

























- in the Adirondack Mountains (New York). *J. Paleolimnol.* 11:189-200.
- Peréz L, Lorenschat J, Massafiero J, Pailles Ch, Sylvestre F, Hollwedel W, Brandorff GO, Brenner M, Islebe G, Lozano MS, Scharf B, Schwalb A, 2013. Bioindicators of climate and trophic state in lowland and highland aquatic ecosystems of the Northern Neotropics. *Rev. Biol. Trop.* 61:603-644.
- Rajapaksa R, Fernando CH, 1987. Redescription and assignment of *Alona globulosa* Daday, 1898 to a new genus *Notoalona* and a description of *Notoalona freyi* sp.nov. *Hydrobiologia* 144:131-153.
- Rey J, Vasquez E, 1986. [Contribution à la connaissance des Cladocères néotropicaux: redescription de *Leydigiopsis ornata* Daday, 1905. (Crustacea, Cladocera)]. [Article in French]. *Ann. Limnol.* 22:169-176.
- Rybak JI, Błędzik LA, 2016. Freshwater Crustacean zooplankton of Europe. Springer: 918 pp.
- Schmidt R, Müller J, Drescher-Schneider R, Krisai R, Szeroczyńska K, Barić A, 2000. Changes in lake level and trophy at Lake Vrana, a large karstic lake on the Island of Cres (Croatia), with respect to palaeoclimate and anthropogenic impacts during the last approx. 16,000 years. *J. Limnol.* 59:113-130.
- Shurin JB, 2000. Dispersal limitation, invasion resistance, and the structure of pond zooplankton communities. *Ecology* 81:3074-3086.
- Sinev AY, 1998. *Alona ossiani* sp. n., a new species of the *Alona affinis* complex from Brazil, deriving from the collection of G.O. Sars (Anomopoda Chydoridae). *Arthropoda Sel.* 7:103-110.
- Sinev AY, 2001. Redescription of *Alona glabra* Sars, 1901. South American species of the pulchella group (Branchiopoda: Anomopoda: Chydoridae). *Arthropoda Sel.* 10:273-280.
- Sinev A, Zawisza E, 2013. Comments on cladocerans of crater lakes of the Nevado de Toluca Volcano (Central Mexico), with the description of a new species, *Alona manueli* sp. Nov. *Zootaxa* 3647:390-400.
- Sinev A Y, 2015a. Revision of the pulchella-group of *Alona s. lato* leads to its translocation to *Ovalona* Van Damme et Dumont, 2008 (Branchiopoda: Anomopoda: Chydoridae). *Zootaxa* 4044:451-492.
- Sinev AY, 2015b. Morphology and phylogenetic position of three species of genus *Camptocercus* Baird, 1843 (Cladocera: Anomopoda: Chydoridae). *Zootaxa* 4040:169-186.
- Sinev AY, Dumont HG, 2016. Revision of the costata-group of *Alona s. lato* (Cladocera: Anomopoda: Chydoridae) confirms its generic status. *Eur. J. Taxon.* 223:1-38.
- Sinev AY, Silva-Briano M, 2012. Cladocerans of genus *Alona* Baird, 1843 (Cladocera: Anomopoda: Chydoridae) and related genera from Aguascalientes State, Mexico. *Zootaxa.* 3569:1-24.
- Sweetman JN, Rühland KM, Smol JP, 2010. Environmental and spatial factors influencing the distribution of cladocerans in lakes across the central Canadian Arctic treeline region. *J. Limnol.* 69:76-87.
- Szeroczyńska K, 1991. Impact of prehistoric settlements on the Cladocera in the sediments of Lakes Suszek, Błędowo, and Skrzetuszewskie. *Hydrobiologia* 225:105-114.
- Szeroczyńska K, Tatur A, Weckstrom J, Gąsiorowski M, Noryskiewicz A, Sienkiewicz E, 2007. Holocene environmental history in northwest Finnish Lapland reflected in the multi-proxy record of small subarctic lake. *J. Paleolimn.* 38:25-47.
- Szeroczyńska K, Zawisza E, 2011a. Subfossil faunal and floral remains (Cladocera, Pediastrum) in two northern Lobelia lakes in Finland. *Knowl. Manag. Aquat. Ect.* 403:09.
- Szeroczyńska K, Zawisza E, 2011b. Records of the 8200 cal BP cold event reflected in the composition of subfossil Cladocera in the sediments of three lakes in Poland. *Quat. Int.* 233:185-193.
- Szeroczyńska K, Zawisza E, Wojewódka M, 2015. Initial time of two high altitude crater lakes (Nevado De Toluca, Central Mexico) recorded in (subfossil) Cladocera. *Stud. Quatern.* 32:109-116.
- Taylor MA, Alfaro E, 2005. Climate of Central America and the Caribbean. 181-189. In: J.E. Oliver (ed.), *Encyclopedia of world climatology*. Springer, Dordrecht.
- ter Braak CJF, 1986. Canonical correspondence analysis: a new eigen vector technique for multivariate direct gradient analysis. *Ecology* 67:1167-1179.
- Van Damme KV, Dumont HJ, 2010. Cladocera of the Lençóis Maranhenses (NE - Brazil): faunal composition and a reappraisal of Sars' method. *Braz. J. Biol.* 70:755-779.
- Van Damme KV, Sinev AY, Dumont HJ, 2011. Separation of *Anthalona* gen.n. from *Alona* Baird, 1843 (Branchiopoda: Cladocera: Anomopoda): morphology and evolution of scraping stenothermic alonines. *Zootaxa* 2875:1-64.
- Zawiska I, Słowiński M, Correa-Metrio A, Obremska M, Luoto T, Nevalainen L, Woszczyk M, Milecka K, 2015. The response of a shallow lake and its catchment to Late Glacial climate changes - A case study from eastern Poland. *Catena* 126:1-10.
- Zawisza E, Szeroczyńska K, 2011. Cladocera species composition in lakes in the area of the Hornsund Fjord (Southern Spitsbergen)-Preliminary results. *Knowl. Manag. Aquat. Ec.* 402:04.
- Zawisza E, Caballero M, Ruiz-Fernandez C, 2012. 500 years of ecological changes recorded in subfossil Cladocera in a high-altitude tropical lake Lago de La Luna, Central Mexico. *Stud. Quatern.* 29:23-29.
- Zawisza E, Correa-Metrio A, Caballero M, Lozano S, Szeroczyńska K, 2014. Paleoecology of tropical Lake Zirhauen (Western Mexico) recorded in Cladocera remains. Abstract book of the 8th Shallow lakes Conf., Antalya, Turkey.
- Zawisza E, Cuna E, Caballero M, Ruiz-Fernandez AC, Szeroczyńska K, Woszczyk M, Zawiska I, 2016. Environmental changes during the last millennium recorded in subfossil Cladocera, diatoms and sediment geochemistry from Lake El Sol (Central Mexico). *Geol. Q.* (in press).