

Revisiting the peculiar regional distribution of muscle fiber types in rat Sternomastoid Muscle

Barbara Ravara (1,2,3), Valerio Gobbo (4), Damiana Incendi (5), Andrea Porzionato (2,5), Veronica Macchi (5), Raffaele De Caro (5), Dario Coletti (6,7,8), Tiziana Martinello (2,9), Marco Patrino (2,9)

(1) *Laboratory of Translational Myology, Department of Biomedical Sciences, University of Padova, Italy;* (2) *Interdepartmental Research Center of Myology (CIR-Myo), University of Padova, Italy;* (3) *A&C M-C Foundation for Translational Myology, Padova, Italy;* (4) *CNR Institute of Neuroscience, Padova, Italy;* (5) *Human Anatomy Section of the Department of Neurosciences, University of Padova, Italy ;* (6) *Sorbonne Universités, UPMC Univ Paris 06 (CNRS, UMR 8256, INSERM ERL U1164), Institut Biologie Paris-Seine, Paris, France;* (7) *Department of Anatomy, Histology, Forensic Medicine & Orthopedics, School of Medicine Sapienza University of Rome, Italy;* (8) *Interuniversity Institute of Myology, Italy;* (9) *Department of Comparative Biomedicine and Food Science, University of Padova, Italy.*

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

The sternomastoid (SM) muscle in rodents is known to have a peculiar distribution of fiber types with a steep gradient from surface to deep region. We here further characterize this peculiar regional distribution by quantitative histochemical morphometries. In Hematoxylin-Eosin (H-E) stained transverse cryosections harvested in the medial portion of the muscle we counted around 10.000 myofibers with a mean diameter of 51.3 ± 12.6 (μm). Cryosections of the SM stained by SDH reaction clearly show two distinct regions, toward the deep surface of the muscle a 40% area that contains packed SDH-positive myofibers, while the remaining area of the SM toward the external surface presents a more checker-board appearance. On the other hand, in the deep region of SM type 1 (slow contracting) muscle fibers, characterized by positive acidic ATPase pH 4.35 reaction, are only the 24.5% of the fibers in the deep area of SM muscles, being restricted to the deepest region. The 75.5% of the myofibers in the deep region are of the fast contracting types (either 48.4% 2A, SDH-positive fibers or 27.1% 2B, SDH-negative fibers, respectively). As expected the 2B muscle fibers, acidic ATPase pH 4.3-negative and SDH-negative, present the largest size, while Type 1 fibers, acidic ATPase pH 4.3-positive and SDH-positive, present the smallest size in rat SM muscle. Based on present and previous observations, comparison of change in absolute number and/or percentage of the fiber types in any experimental model of muscle atrophy/hypertrophy/plasticity/pathology /recovery in the rat SM, and possibly of all mammals, will ask for morphometry of the whole muscle cross-sections, muscle sampling by bioptic approaches will provide only comparable data on the size of the different types of muscle fibers.

Key Words: rat, Sternomastoid muscle, muscle fiber types, SDH, myofibrillar ATPases, regional distribution

Eur J Transl Myol 28 (1): xxx-xxx, 2018

The sternomastoid muscle (SM) in rodents is known to have a peculiar distribution of fiber types with a steep gradient from surface to deep region.¹ This behavior is also present in leg muscles (e.g., tibialis anterior muscle), but not at extent of the SM muscle.²⁻⁴ We are characterizing in normal animals this peculiar regional distribution by quantitative histochemical analyses of

transverse sections in the third medial portion of the muscle. We will present results suggesting that comparison of change in absolute number and/or percentage of the fiber types in the rat SM will ask for morphometry of the whole muscle cross-section, while muscle bioptic samplings will provide only consistent results on size of muscle fibers.

Regional distribution of muscle fiber types in rat sternomastoid muscle

Eur J Transl Myol 28 (1): 117-123, 2018




Fig 1. Transverse cross-section of normal rat SM muscle. Panel A, SDH reaction. In the deep region are present small diameter, intense blue fibers, while the superficial region presents higher diameter unstained fibers (type 2B) and some light to middle blue fibers (Type 2A). Panel B, ATPase pH 4.35 reaction. The black fibers (type 1) are restricted to deepest region of the SM muscle. Scale Bar 1 mm.

Materials and Methods

Animals and harvesting of muscles

Adult females Wistar rats, 300 g of weight were used. These animals come from the vivarium of the Department of Neurosciences – University of Padova.

The animals were kept in cages with light/dark cycle of 12h, room temperature (24 ± 2), food and water *ad libitum*. All procedures in this study were according to the international ethical principles of animal experimentation and approved by the Ethics on Animal Experimentation Committee (EAEC) of University of

Regional distribution of muscle fiber types in rat sternomastoid muscle

Eur J Transl Myol 28 (1): 117-123, 2018

Table 1. Mean myofiber diameter (μm) in the SM rat muscle stained by H-E. Values are mean \pm SD

All muscle fibers	SM dx	SM sx
	51.6 \pm 26.7	50.9 \pm 12.1
Mean of two SM	51.3 \pm 12.7	
Total fibers in Deep and Superficial region	9981	

Padova. Animals were killed by deep anaesthesia and muscles were excised, weighted, frozen in liquid nitrogen (-196°C) and kept at -80°C until use.

Histology and histochemistry of the sternomastoid muscle

Size, and absolute or percent contents of different muscle fiber types were determined in transverse cryosections of the SM muscle harvested in the medial third of the muscle. Serial $10\mu\text{m}$ cross-sections were cut on a cryostat microtome at -25°C , mounted on slides and stained in small glass jars. From stained sections digital images were collected at magnification of $\times 2.5$ and $\times 10$ using a transmission light microscope mounting a HDMI camera. Quantitative evaluations were performed on the collected pictures using Scion Image for Windows version Beta 4.0.2 (2000 Scion Corporation). Muscle fiber sizes and regional distribution of the muscle fiber type were determined using the Adobe Photoshop software (Adobe Systems Incorporated, San Jose, CA).

Absolute number of muscle fibers in rat SM muscle

To determine the absolute number of muscle fibers in the rat SM muscle, cryosections were stained with standard Hematoxylin-Eosin (H-E) procedure.⁵

Fiber typing of the muscle fibers was performed by the following histochemical methods

Succinate Dehydrogenase (SDH) reaction

SDH reaction was performed to distinguish between oxidative, "less" oxidative and non-oxidative muscle fibers. Muscle sections were incubated in SDH incubation solution for 60 min at 37°C . Sections were rinsed in distilled H_2O , 3×1 min and removed unbound NBT from sections with 3×1 min exchanges of acetone solutions 30, 60, 90 % in distilled H_2O , in increasing then decreasing concentrations. Sections were rinsed in distilled H_2O , 3×1 min and dehydrated in graded alcohol solutions (ethanol 70, 90, 100%) cleared in xylene and mounted in permanent medium: Canada balsam. SDH incubating solution: NBT (nitro blue tetrazolium): 10mg; SDH stock solution: 10ml. Adjust to pH 7.2 to 7.6 with 0.1 N HCl or NaOH. SDH stock solution: 0.2M sodium

succinate solution 100ml, 0.2M Phosphate buffer, pH 7.4 100ml, make freshly at least every two weeks.

Histochemistry by Myofibrillar actomyosin ATPase

Two different procedures were used for staining myofibrillar actomyosin ATPase using essentially the same methods described by Brooke and Kaiser,^{6,7} and by Guth and Samaha.⁸⁻¹⁰ For determination of the alkali preincubated myofibrillar ATPase activity the following procedures were employed. cryosections were air dried for 10 min at room temperature and fixed for 3 min in 5% (w/v) formaldehyde in 200mM sodium cacodylate, 68mM $\text{CaCl}_2 \cdot 2\text{H}_2\text{O}$ and 340 mM sucrose (adjusted with HCl to pH 7.6 before adding the formaldehyde). After washing in wash solution 100mM Tris, 18mM CaCl_2 (pH 7.8), an alkaline pre-incubation was performed for 15 min at room temperature in a solution containing 200mM 2-amino-3-methyl-1 propanol (adjusted with HCl to pH 10.4). After two successive 1 min incubation in the wash solution, sections were incubated for 45 min at 37°C in ATP solution (2,7mM ATP, 90mM CaCl_2 , 100mM sodium barbital adjusted with NaOH to pH 9.4). For the acid preincubation myofibrillar ATPase activity (10, 11), the following procedures were utilized. Sections were air dried for 10 min at room temperature and incubated in a solution consisting of 100mM sodium acetate (adjusted with HCl to pH 4.35) for 10 min at room temperature. After washing 2×1 min in 18 mM CaCl_2 and 100mM Tris HCl pH 7.8, the sections were incubated at 37°C for 45 min in ATP solution together to serial sections with alkali preincubation. After washing 2×30 sec in distilled H_2O , sections were incubated for 5 min in 2% (w/v) CoCl_2 , washed 2×30 sec in distilled H_2O , sections were incubated 5 min in 2% (v/v) $(\text{NH}_4)_2\text{S}$, washed in distilled H_2O 2×30 sec, and dehydrated in graded alcohol solution (ethanol 70-90-100%) cleared in xylene, and mounted with Canada balsam medium.

Results

Morphometry: number and size of rat SM muscle fibers in H-E stained cryosections.

In the rat SM muscle cryosections harvested in the medial portion of the muscle we counted around 10.000

Regional distribution of muscle fiber types in rat sternomastoid muscle

Eur J Transl Myol 28 (1): 117-123, 2018

myofibers. The mean diameter in these muscle sections stained by H-E was $51.3 \pm 12.7 \mu\text{m}$ (Table 1).

Histochemistry of sternomastoid rat muscle

Fiber typing by SDH reaction

In Fig.1A is presented a transverse section from the medial region of a rat SM muscle stained by SDH reaction. The cryosection clearly shows two distinct regions, a right part of the section that cover around 40% area that contains SDH positive myofibers (right part of Fig.1A) and a the left part of Fig.1A; the latter encompasses the other 60% area of the SM with a more checker-board appearance. The right part of the cryosection, peculiarly rich in SDH positive muscle fibers, corresponds to the deep part of the SM, while the left part of the figure correspond to the superficial region of the SM. Table 2 shows that SDH-positive myofibers are the 72.9% of the total myofibers of SM deep region. In the superficial region (left part of the cryosection) based on SDH staining (Fig.1A), the mitochondrial rich, 2A muscle fibers are 31.1%. Since very few or none of type 1 muscle fibers are present there (see ATPase histochemistry reaction). The 68.9% 2B negative fibers are that of the myofibers present in this region of the SM muscle. Figure 2 shows six representative fields at higher magnification taken from periferal, intermediate, and deep portions of cryosections stained either by SDH (A, B, C) or acidic ATPase (D, E, F).

Histochemistry by Myofibrillar actomyosin ATPase

In the deep region (right part of Fig.1B), the type 1 (slow contracting) muscle fibers, identified by positive acidic ATPase pH 4.35 (Fig.1B), are only the 24.5% of the fiber of this part of SM muscle (Table2). On the other hand, the slow type muscle fibers, i.e., those positive at the

acidic ATPase pH 4.35 are restricted in the deepest region of SM muscle. Thus, 75.5% of the myofibers in the deep region are of the fast contracting types (either 48.4% 2A and 27.1% are 2B, respectively).

In the superficial region, left part of the cryosection, based on acidic ATPase pH 4.35, the type 1 positive fibers are absent, the type 2A and muscle fibers characterized by an intermediate intensity decrease to 20.3%. Therefore the vast majority of the muscle fibers present in the superficial region of the SM muscle are of the 2B type (68.9% as SDH- negative fibers or 79.7% when counted in the acidic ATPase pH 4.35).

Fiber size of the three different fiber types of the SM muscle

As expected, Table 1, shows that the acidic ATPase pH 4.35 negative and SDH-negative 2B type myofibers, present the largest size among the three types of muscle fiber of rat SM muscle. In the superficial region based on acid ATPase pH 4.35, the 2B fiber size result $47.6 \mu\text{m}$ and the 2A fiber $35.3 \mu\text{m}$. In the deep region the fiber size of 1 fibers (slow contracting) result to be the smallest: $29.2 \mu\text{m}$; while the 2A and 2B fiber size are $36.8 \mu\text{m}$ on average in size. Based on mitochondrial enzyme activities, SDH reaction, in the superficial region the size of 2B fiber recognizable as the larger ones are $53.3 \mu\text{m}$ and in the deep region $41.1 \mu\text{m}$, while the 1 and 2A fiber size in the superficial region and in the deep region is respectively $35.8 \mu\text{m}$ and $33.8 \mu\text{m}$.

Discussion

The peculiarity of the SM rat muscle is the presence of two distinct regions, where the fiber type composition is characterized in both zone by 2B fibers that are easily

Table 2. Mean myofiber diameter (μm) in deep and superficial region of rat SM stained by SDH or ATPase, pH 4.35 reactions. Values are mean \pm SD.

SDH reaction			
Deep region		Superficial region	
Type fibers 1 + 2A	33.8 ± 7.5	35.8 ± 9.2	
Type fibers 2B	41.1 ± 9.2	53.3 ± 10.8	
Total number fibers	5211	3975	
ATPase pH 4.35			
Deep region		Superficial region	
Type fibers 1	29.2 ± 6.5	Type fibers 2A	35.3 ± 8.5
Type fibers 2A + 2B	36.8 ± 9.2	Type fibers 2B	47.6 ± 11.9
Total number fibers	5802	4179	

Regional distribution of muscle fiber types in rat sternomastoid muscle

Eur J Transl Myol 28 (1): 117-123, 2018

Table 3 . Number and percentage of SM fiber types in deep or superficial regions by SDH or ATPase pH 4.3

SDH reaction			
	Deep region		Superficial region
Type fibers 1+2A			
Number	3800		1235
Percentage	72.9%		31.1%
Type fibers 2B			
Number	1411		2740
Percentage	27.1%		68.9%
Total number fibers	5211		3975
Total fibers Deep plus Superficial region	9186		
ATPase pH 4.35			
	Deep region		Superficial region
Type fibers 1		Type fibers 2A	
Number	1422		849
Percentage	24.5%		20.3%
Type fibers 2A+ 2B		Type fibers 2B	
Number	4380		3330
Percentage	75.5%		79.7%
Total number fibers	5802		4179
Total fibers Deep and Superficial region	9981		

recognizable from their major size, while the 1 and 2A are the smallest. The SM muscle presents a very heterogeneous distribution of muscle fiber types with a large predominance of the type 2B in the superficial region while moving to deep regions there is a progressive enrichment of 2A fibers. Only in the deepest region of the SM muscle there are type 1 muscle fibers (Fig 1B and Table 3).

It remains to be determined if the content of fiber types varies along the muscle length and if applying new imaging processing as the Machine-learning algorithms (Deep Neural Networks) that have been proved to be very powerful methods for automatic image segmentation, especially in the field of histological analyses,¹¹ some of the problems we are facing with non homogeneous muscles would be overcome. Anyhow, as demonstrated here and by previously,¹ in the rat SM muscle comparative analyses for experimental models of muscle plasticity, in normal and diseased muscles,¹²⁻²⁶ will require morphometry of the whole cross-sections. In conclusion, in case of bioptic fragments of SM muscle, in small, but conceivably also large, mammals, only the size of the different fiber types could be evaluated to

avoid systematic errors related to different regional sampling.

List of acronyms

ATPase - myofibrillar Actomyosin ATPase histochemistry
EAEC - Ethics Committee on Animal Experimentation
H-E - Hematoxylin-Eosin reaction
SDH - succinate dehydrogenase reaction
SM - sternomastoid muscle

Author's contributions

Authors equally contributed to the manuscript.

Acknowledgments and Funding

BR thanks for support A&C M-C Foundation for Translational Myology, Padova, Italy.

Conflict of Interest

The authors declare no conflicts of interests.

Ethical Publication Statement

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this

typescript is consistent with those guidelines.

Corresponding Author

Barbara Ravara, Laboratory of Translational Myology, of the Interdepartmental Research Center for Myology (CIR-Myo), c/o Department of Biomedical Sciences, University of Padova and A&C M-C Foundation for Translational Myology, Padova, Italy.

E-mail: barbara.ravara@unipd.it

E-mails of co-authors

Damiana Incendi: damiana.incendi@unipd.it

Andrea Porzionato: andrea.porzionato@unipd.it

Valerio Gobbo: gobbov@bio.unipd.it

Veronica Macchi: veronica.macchi@unipd.it

Raffaele De Caro: Raffaele.decar@unipd.it

Dario Coletti: dario.coletti@snv.jussieu.fr

or dario.coletti@uniroma1.it

Tiziana Martinello: tiziana.martinello@unipd.it

Marco Patruno: marco.pat@unipd.it

References

- Polican Ciena A, Yokomito de Almeida SR, de Matos Alves PH, et al. Histological and ultrastructural changes of sternomastoid muscle in aged wistar rats. *Micron* 2011;42:871-6.
- Hiroux C, Vandoorne T, Koppo K, et al. Physical Activity Counteracts Tumor Cell Growth in Colon Carcinoma C26-Injected Muscles: An Interim Report. *Eur J Transl Myol* 2016;26:5958. doi: 10.4081/ejtm.2016.5958. eCollection 2016 Jun 13.
- Tasic D, Dimov D, Gligorijevic J, et al. Muscle fibre types and fibre morphometry in the tibialis posterior and anterior of the rat: a comparative study. *Medicine and Biology* 2003;10:16-21.
- Coletti D, Daou N, Hassani M, et al. Serum Response Factor in Muscle Tissues: From Development to Ageing. *Eur J Transl Myol* 2016;26:6008. doi: 10.4081/ejtm.2016.6008. eCollection 2016 Jun 13.
- Kern H, Loeffler S, Hofer C, et al. FES Training in Aging: interim results show statistically significant improvements in mobility and muscle fiber size. *Eur J Transl Myology* 2012;22:61-7.
- Brooke MH, Kaiser KK. J. Some comments on the histochemical characterization of muscle adenosin triphosphatase. *Histochem Cytochem* 1969;17:431-2.
- Brooke MH, Kaiser KK. Muscle fiber types: How many and what Kind? *Arch Neurol* 1970 23; 369-79.
- Guth L, Samaha FJ. Qualitative differences between actomyosin ATPase of slow and fast mammalian muscles *Exp Neurol* 1969;25:139-52.
- Guth L, Samaha FJ. Procedure for the histochemical demonstration of actomyosin ATPase. *Exp Neurol* 1970;28: 365-7.
- Hammalinen N, Pette D. The histochemical profiles of fast fibers IIB, IIA in skeletal muscle of mouse, rat and rabbit. *Histochem Cytochem* 1993;41:733-43.
- Ciresan DC, Giusti A, Gambardella LM, Schmidhuber J. Mitosis detection in breast cancer histology images with deep neural networks. *Med Image Comput Assist Interv* 2013;16:411-8.
- Seene T, Umnova M, Kaasik P. Morphological peculiarities of neuromuscular junctions among different fiber types: Effect of exercise. *Eur J Transl Myol*. 2017 Jun 27;27(3):6708. doi: 10.4081/ejtm.2017.6708. eCollection 2017 Jun 27.
- Samiee F, Zarrindast MR. Effect of electrical stimulation on motor nerve regeneration in sciatic nerve ligated-mice. *Eur J Transl Myol*. 2017 Sep 20;27(3):6488. doi: 10.4081/ejtm.2017.6488. eCollection 2017 Jun 27.
- Power GA, Dalton BH, Gilmore KJ, et al. Maintaining Motor Units into Old Age: Running the Final Common Pathway. *Eur J Transl Myol* 2017;27(1):6597. doi: 10.4081/ejtm.2017.6597. eCollection 2017 Feb 24. Pigna E, Greco E, Morozzi G, et al. Denervation does not Induce Muscle Atrophy Through Oxidative Stress. *Eur J Transl Myol* 2017;27(1):6406. doi: 10.4081/ejtm.2017.6406. eCollection 2017 Feb 24.
- Pette D, Vrbová G. The Contribution of Neuromuscular Stimulation in Elucidating Muscle Plasticity Revisited. *Eur J Transl Myol*. 2017 Feb 24;27(1):6368. doi: 10.4081/ejtm.2017.6368. eCollection 2017 Feb 24. Review.
- Carotenuto F, Coletti D, Di Nardo P, Teodori L. α -Linolenic Acid Reduces TNF-Induced Apoptosis in C2C12 Myoblasts by Regulating Expression of Apoptotic Proteins. *Eur J Transl Myol* 2016;26(4):6033. doi: 10.4081/ejtm.2016.6033. eCollection 2016 Sep 15.22.
- Riebold B, Nahrstaedt H, Schultheiss C, et al. Multisensor Classification System for Triggering FES in Order to Support Voluntary Swallowing. *Eur J Transl Myol* 2016;26(4):6224. doi: 10.4081/ejtm.2016.6224. eCollection 2016 Sep 15.33.
- Stratton K, Faghri PD. Electrically and Hybrid-Induced Muscle Activations: Effects of Muscle Size and Fiber Type. *Eur J Transl Myol* 2016;26(3):6163. eCollection 2016 Jun 13.
- Willand MP. Electrical Stimulation Enhances Reinnervation After Nerve Injury. *Eur J Transl Myol* 2015;25(4):243-8. doi: 10.4081/ejtm.2015.5243. eCollection 2015 Aug 24. Review.
- Zampieri S, Mosole S, Löffler S, et al. Physical Exercise in Aging: Nine Weeks of Leg Press or Electrical Stimulation Training in 70 Years Old Sedentary Elderly People. *Eur J Transl Myol*

Regional distribution of muscle fiber types in rat sternomastoid muscle

Eur J Transl Myol 28 (1): 117-123, 2018

- 2015;25(4):237-42. doi: 10.4081/ejtm.2015.5374. eCollection 2015 Aug 24. Review.
21. Barber L, Scicchitano BM, Musaro A. Molecular and Cellular Mechanisms of Muscle Aging and Sarcopenia and Effects of Electrical Stimulation in Seniors. *Eur J Transl Myol* 2015;25(4):231-6. doi: 10.4081/ejtm.2015.5227. eCollection 2015 Aug 24. Review.
 22. Ravara B, Gobbo V, Carraro U, et al. Functional Electrical Stimulation as a Safe and Effective Treatment for Equine Epaxial Muscle Spasms: Clinical Evaluations and Histochemical Morphometry of Mitochondria in Muscle Biopsies. *Eur J Transl Myol* 2015;25(2):4910. doi: 10.4081/ejtm.2015.4910. eCollection 2015 Mar 11.
 23. Costa A, Rossi E, Scicchitano BM, et al. Neurohypophyseal Hormones: Novel Actors of Striated Muscle Development and Homeostasis. *Eur J Transl Myol* 2014;24(3):3790. doi: 10.4081/ejtm.2014.3790. eCollection 2014 Sep 23. Review.
 24. Hockerman GH, Dethrow NM, Hameed S, et al. The Ubr2 Gene is Expressed in Skeletal Muscle Atrophying as a Result of Hind Limb Suspension, but not Merg1a Expression Alone. *Eur J Transl Myol*. 2014;24(3):3319. doi: 10.4081/ejtm.2014.3319. eCollection 2014 Sep 23.
 25. Lomo T. The Response of Denervated Muscle to Long-Term Stimulation (1985, Revisited here in 2014). *Eur J Transl Myol* 2014;24:3294. doi: 10.4081/ejtm.2014.3294. eCollection 2014 Mar 31.
 26. Carlson BM. The Biology of Long-Term Denervated Skeletal Muscle. *Eur J Transl Myol*. 2014 Mar 27;24:3293. doi: 10.4081/ejtm.2014.3293. eCollection 2014 Mar 31.

Received for publication: January 15, 2018

Revision received: February 13, 2018

Accepted for publication: February 14, 2018