

## SUPPLEMENTARY MATERIAL

### The impact of glomerular filtration rate on short-term outcome in elderly patients with heart failure with preserved ejection fraction

Giuseppe Delli Carpini, Cristina Di Bartolomeo, Gianluca Testa, Germano Guerra, Klara Komici

Department of Medicine and Health Sciences, University of Molise, Campobasso, Italy

**Correspondence:** Klara Komici, Department of Medicine and Health Sciences, University of Molise, Campobasso, 86100 Italy.

E-mail: klara.komici@unimol.it

**Key words:** heart failure, heart failure with preserved ejection fraction; chronic kidney disease.

**Supplementary Table 1. Logistic regression analysis.**

Variable	Odds ratio	95% confidence interval	p
Age	1.08	1.01-1.15	0.013
Gender	0.28	0.097-0.82	0.02
DM	0.67	0.21-2.22	0.52
AF	1.19	0.42-3.33	0.74
ACEi/ARB	1.04	0.31-3.45	0.94
Aldosterone antagonists	3.64	1.0-12.3	0.08
$\beta$ -blockers	1.00	0.35-2.87	0.99
WBC	1.01	0.91-1.14	0.74
Haemoglobin	1.14	0.86-1.53	0.34
eGFR	0.95	0.91-0.98	0.009

DM, diabetes mellitus; AF, atrial fibrillation; ACE-i/ARB, angiotensin converting enzyme inhibitors/angiotensin receptor blockers; WBC, white blood cells; eGFR, estimated glomerular filtration rate