



eISSN: 2281-7824

<https://www.pagepressjournals.org/index.php/hls/index>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The **Early Access** service lets users access peer-reviewed articles well before print / regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Healthcare in Low-resource Settings** is, therefore, e-publishing PDF files of an early version of manuscripts that undergone a regular peer review and have been accepted for publication, but have not been through the typesetting, pagination and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear on a regular issue of the journal.

E-publishing of this PDF file has been approved by the authors.

Healthc Low-resour S 2024 [Online ahead of print]

To cite this Article:

Abdul Majeed AA, Haleem AA. **Prevalence of hypertension and its associated risk factors among secondary school students in Duhok City.** Healthc Low-resour S doi: 10.4081/hls.2024.12073

 ©The Author(s), 2024

Licensee [PAGEPress](#), Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Prevalence of hypertension and its associated risk factors among secondary school students in Duhok City

Ayoub Abid Abdul Majeed,¹ Azad AbdulJabar Haleem²

¹Master in Pediatric Nursing, Psychiatry and Pediatric Nursing Unit, College of Nursing, University of Duhok; ²Pediatric Department, College of Medicine, University of Duhok; Heevi Pediatrics Teaching Hospital; Kurdistan Pediatrics Society, Iraq

Correspondence: Ayoub Abid Abdul Majeed, Psychiatry and Pediatric Nursing Unit, College of Nursing, University of Duhok, Iraqi Kurdistan.

E-mail: ayoub.abid@uod.ac

Key words: cross-sectional studies; hypertension; prevalence; risk factors.

Conflict of interest: the authors declare no potential conflict of interest, and all authors confirm accuracy.

Funding: We did not have any funding to conduct this study.

Ethics approval: the Ethics Committee of Duhok General Directorate of Health approved this study (registered as 1307202I-7 -2I on 13 Jul,2021. The study is conformed with the Helsinki Declaration of 1964, as revised in 2013, concerning human and animal rights.

Informed consent: The written consent forms were not applicable in this study because we collected the data of this study from the students, not patients

Patient consent for publication: Written informed consent was obtained from a legally authorized representative(s) for anonymized Students information to be published in this article.

Availability of data and materials: all data generated or analyzed during this study are included in this published article.

Acknowledgments: We would like to present our deep thanks to the Education of Duhok, the secondary and high schools, and the students who participated in this study.

Abstract

Hypertension can develop in early childhood and go unnoticed unless it is screened for specifically during this developmental stage. Detecting potential hypertension risk in children is critical to avoiding the serious, long-term complications associated with the condition. The purpose of this study was to investigate the prevalence of hypertension among schoolchildren aged 13 to 18, as well as the risk factors associated with it. A cross-sectional study included 565 students aged 13 to 18 from the Duhok region. The analysis used Chi square testing and logistic regression with JMP Pro 14.3.0 software. Of the 565 students from 32 schools chosen, 242 (42.80%) were male and 323 (57.2%) were female. Notably, 5.84% (n=33) were hypertensive. A significant correlation was found between hypertension prevalence and body mass index, as well as gender. There was no significant relationship found with other variables. Logistic regression revealed that high blood pressure was significantly associated with excess weight, obesity, and age. We concluded that there was an increase in the incidence of hypertension among high school students. Lifestyle changes appeared to play a role in hypertension development among this population. Routine school surveys are recommended to

detect potential hypertension cases in children and adolescents, allowing for timely preventative interventions.

Introduction

Hypertension poses a significant global health challenge and stands as a primary contributor to coronary artery and cerebrovascular diseases. The year 2000 witnessed around one billion adults worldwide grappling with hypertension, a number anticipated to escalate to 1.56 billion by 2025.^{1,2} This pervasive ailment entails substantial morbidity and mortality, silently endangering populations globally. Evidently, hypertension can trace its roots back to childhood, remaining concealed unless diligently sought out during this developmental phase. Thus, timely identification of hypertension and its triggers assumes paramount importance to forestall its complications.³

Given the paucity of symptoms during childhood and adolescence, hypertension often evades detection during these formative years. Its prevalence escalates with age and adolescence, rooted in multifaceted factors such as genetics, race, geography, culture, and dietary habits.³ Intriguingly, elevated blood pressure in this period foreshadows similar levels in adulthood and old age. Our understanding of blood pressure in children and adolescents is far from being complete and the long-term natural history of blood pressure in this age group is still not well understood.⁴

Hypertension seldom manifests symptoms during early life, though the WHO does not advocate hypertension screening for children and adolescents. This study was undertaken due to a lack of data on adolescent hypertension prevalence in our country. Hypertension poses a substantial public health challenge among adults globally, with Iraq registering a 40.4% prevalence in a 2006 non-communicable disease risk factor survey.⁵

Insufficient information exists regarding blood pressure patterns and hypertension prevalence in Iraqi children and other developing nations. Notably, a 2009 Najaf City study on school

students found a 5.5% prevalence of high blood pressure while a 2002 Baghdad study on school-aged students reported a 1.7% hypertension prevalence.⁶

Despite strides in hypertension detection and management in select countries, the global scenario concerning awareness, treatment, and control remains wanting.¹ To ensure public awareness and regulation of hypertension, early identification in life's initial stages becomes pivotal. Delayed recognition burdens not only individuals but also governments due to elevated treatment costs. This study seeks to unveil hypertension prevalence and associated risk factors among secondary school students in the Kurdistan Region, thereby contributing to our understanding of this critical health issue.

Materials and Methods

Study design

The study was carried out in secondary and high schools in Duhok provinces in the Kurdistan region (north of Iraq). A cross-sectional study design was used; the study was carried out from 3rd November 2021 to 10th April 2022. Ethical approval from the Directorate General of Health/ Duhok and Directorate General of Education/ Duhok was obtained. Consent was obtained from the respondents before the interview with their parent's consent.

Sampling technique

In this research, the target group comprised adolescents aged 13 to 18, encompassing both genders and attending school. It was crucial for respondents to be an accurate representation of the intended population. To ensure this, a comprehensive multistage sampling technique was employed. In the initial phase, Duhok City was categorized into two distinct zones through stratified sampling. Subsequently, a list of schools in each zone was compiled, from which a total of thirty-two government schools were arbitrarily chosen to adequately represent both male and female secondary education. This selection was carried out using a simple random sampling method, specifically the lottery approach. This resulted in a random yet representative subset extracted from the overall population, allowing for a more precise estimation of characteristics. The sample size, calculated at a 95% confidence level, was established as 565 students. Students who were excluded from participation included those taking study breaks and those whose parents withheld consent. Additionally, students with chronic medical

conditions or those reliant on specific medications affecting blood pressure, like steroids, were also omitted from the study.

Instruments and data collection

The collection of data will involve the utilization of a questionnaire, meticulously crafted through an exhaustive analysis of preceding research and literature pertinent to the current study's subject matter. This questionnaire comprises two distinct segments for data compilation. The first section of the questionnaire will be administered through face-to-face interviews with the participants. The second part, on the other hand, will involve data collection from the parents of the participants.

Statistical methods

The demographic information of school children was depicted using numerical values and percentages. The occurrence rate of hypertension in school children was ascertained both in numerical counts and percentages. This was achieved by dividing the number of patients exhibiting various degrees of hypertension by the total count of school children and then multiplying by 100. The assessment of hypertension prevalence based on factors like age, gender, and other general characteristics was conducted through the utilization of the Pearson chi-squared test. Similarly, the connection between hypertension in school children and parental socio-demographic traits and their lifestyles was evaluated using the Pearson chi-squared test. To identify predictors of hypertension among school children, nominal logistic regression was employed. A p-value below 0.05 was deemed indicative of statistically significant differences. The statistical computations were conducted using JMP Pro 14.3.0 software.

Anthropometric measurements

Anthropometric measurements were conducted within the school setting. Each measurement was taken twice, and the average of the two readings was considered. For height measurement, participants were barefoot and measured using a portable stadiometer (Seca 213, Measuring range 20 - 200cm). Weight was recorded with participants wearing lightweight attire, employing a digital Heine portable scale accurate to the nearest 0.1kg. BMI was computed for each child by dividing their weight (in kg) by the square of their height (in meters) (kg/m^2). To gauge the appropriateness of participants' BMI values, reference data from the most recent anthropometric standards specific to their sex and age were sourced from the Centers for

Disease Control and Prevention (CDC). These reference standards were used to establish percentiles for height, weight, and BMI, aiding in the assessment of participants' BMI in relation to their age and sex.⁷

Children were grouped into categories determined by these percentiles as outlined: Individuals below the 5th percentile were classified as underweight, those between the 5th percentile and less than the 85th percentile were considered to have a healthy weight, those between the 85th percentile and less than the 95th percentile were classified as overweight, and those equal to or surpassing the 95th percentile were categorized as obese. For adults aged 20 years and older (*i.e.*, children's parents), the classification was slightly different. Adults with a BMI ranging from 18.5 to less than 25 were labeled as having a normal weight, those with a BMI between 25 and less than 30 were identified as overweight, and individuals with a BMI of 30 or higher were categorized as obese.⁸

Blood pressure measurement

Blood pressure assessments were conducted using a standardized mercury sphygmomanometer, paired with a manually inflated cuff of appropriate size, and a stethoscope. Measurements were taken from the right arm while the child was in a calm sitting position for a minimum of 5 minutes to alleviate any anxiety. The researcher inquired about recent activities such as smoking, eating, exertion, or exercise within the last 30 minutes before participating in the study.

The student was positioned comfortably, with their back supported, feet resting on the floor, the right arm properly supported, and the cubital fossa (elbow crease) aligned with the level of the heart. If the blood pressure reading equaled or exceeded the 90th percentile based on the child's age, gender, and height, the blood pressure measurement was repeated twice during the same visit. The average values for systolic blood pressure (SBP) and diastolic blood pressure (DBP) were utilized.

If the mean SBP and/or DBP readings were elevated during the initial visit, two additional blood pressure measurements were performed within a two-week timeframe. This multi-step approach ensured accurate and reliable blood pressure assessments, taking into consideration possible variations in measurements.

Results

The study showed that 15.04% of the school children had abnormal blood pressure. The classes of abnormal blood pressure were Elevated blood pressure (7.43%), Hypertensive (5.84%), Stage 1 hypertensive (0.71%), and Stage 2 hypertensive (1.06%). The children were males (42.83%) and females (57.17%). The children had different age groups from 13 to 18. They reside in urban (10.09%) and rural areas (89.91%) and had different family size (Table 1). The study showed that the prevalence of different classes of abnormal blood pressure was not statistically different based on the gender ($p=0.0934$), age groups ($p=0.2338$), residency ($p=0.8967$), and family size ($p=0.7519$; Table 2).

The study showed that the children whose fathers had lower level of education were more likely to have abnormal blood pressures ($p=0.033$). In addition, those children who had the obese fathers were more likely to have different classes of abnormal blood pressures ($p=0.0421$). The similar pattern was found for the children whose fathers had hypertension ($p=0.0345$) and sleep apnea ($p=0.0002$). In terms of mothers, the similar pattern was found for the education and hypertension. Also, the children who had employee mother were more likely to have hypertension (Table 3).

The study showed that the children who consumed the free fat in their food by the family were more likely to be affected by abnormal blood pressure ($p=0.0375$). In addition, the children who were obese were more likely to be affected by abnormal blood pressure compared to those children with normal and underweight ($p<0.0001$; Table 4). The study showed that the age and obesity of the children predicted the blood pressure among secondary school children (Table 5; Figure 1). The study did not show the significant difference of QoL among children with different classes of abnormal blood pressure ($p=0.2103$ Table 5; Figure 2).

Discussion

Although the occurrence of clinical hypertension among children is significantly lower compared to adults, substantial evidence indicates that the origins of essential hypertension can be traced back to childhood. Numerous epidemiological investigations have addressed blood pressure levels in children. However, there exists considerable diversity in the methodologies employed and in the criteria defining normal and elevated blood pressure. Just as in adults, a variety of environmental and genetic factors impact children's blood pressure. As a result, it is crucial to conduct extensive population-based research efforts to gather standardized reference data concerning blood pressure levels in children.

Symptoms arising from hypertension are infrequent during the early stages of life. Importantly, it should be highlighted that the World Health Organization (WHO) does not endorse routine blood pressure screening for children and adolescents.⁹ Much like in adults, the blood pressure of children is impacted by various environmental and genetic factors. Consequently, it is essential to conduct population studies to gather standard reference data concerning blood pressure levels in children. In the early years of life, symptoms stemming from hypertension are uncommon. It's important to note that the World Health Organization (WHO) does not endorse routine blood pressure screening for children and adolescents.¹⁰

Simultaneously, the notion of the cardiovascular continuum underscores the importance of adopting preventive measures at the earliest opportunity. Being aware of changes at a young age can trigger early preventive strategies, yielding favorable long-term outcomes. Our motivation for conducting this prevalence survey is the lack of data on hypertension prevalence among adolescents in our country. Hypertension, a significant contributor to coronary artery disease and stroke, originates in childhood. Screenings for hypertension, even in cases of borderline hypertension and when risk factors are present, can significantly enhance the quality of life for these children, helping them lead healthier lives.¹¹

Prevalence of hypertension

In this current investigation, we present data pertaining to blood pressure readings from a representative group of 565 school children aged 13 to 18 in Duhok. Through a comprehensive analysis involving three distinct visits, each comprising triplicate blood pressure measurements, we identified that the overall prevalence of hypertension among this cohort, spanning ages 13 to 18, was 5.84%. The study sample encompassed individuals aged 13 to 18 years. These findings become particularly concerning when juxtaposed with the outcomes of other investigations, like the study conducted by Macedo *et al.* In their research, they examined the prevalence of hypertension in 889 children aged 5 to 18 in northern Portugal, specifically exploring its correlation with obesity. Their study reported an estimated hypertension prevalence of 5.2%, which is remarkably close to the prevalence observed in our study.¹²

This value aligns well with our own findings concerning prevalence. Furthermore, the occurrence and frequency of hypertension among school children in Turkey have been subjects of investigation by numerous researchers. The hypertension rates have exhibited variations across studies conducted in various countries, including those carried out within Turkey. In the context of school children studied in Shimla, north India, the observed prevalence of

hypertension was 5.9%. This rate of prevalence notably exceeded the predicted 5% prevalence of high blood pressure in children in the United States using the same criteria.¹³

Across the board, systematic reviews have reported varying estimates for hypertension in children, with Africa having a prevalence estimate of 5% and America showing a prevalence of 3.5%.¹⁴ Another study in 2014 by Patel *et al.* from Bhopal documented a prevalence of hypertension at 5.36%. In various regions of Africa, such as Ghana in West Africa and Tunisia in North Africa, prevalence rates of 6% and 9.6%, respectively, were documented by Addo *et al.* and Harrabi *et al.*¹⁵ Similarly, in a study by Kidy *et al.*, the prevalence of hypertension was examined.¹⁶

The prevalence of hypertension reported in this study is consistent with the prevalence reported in the literature. The reason for the difference between the prevalence of hypertension in this study and those in the previous studies might be the usage of different age groups.¹⁷ and maybe because of different food intake and lifestyles in different populations.

Relationship between BMI and hypertension

Blood pressure demonstrated a positive correlation with BMI (height and weight), and this association remained statistically significant ($p=0.00$) according to the results of our multivariate regression analysis. In our study, a notable trend emerged wherein the prevalence of hypertension heightened as BMI status increased, showcasing strong statistical significance. This finding reinforces the conclusions drawn by Jonathan *et al.*, underscoring the substantial risk factors that obesity and overweight represent in relation to hypertension.¹⁸ This linkage, where higher blood pressure levels correspond with increased obesity rates, has been consistently documented in a multitude of reports.¹⁹

This study uncovered a notable pattern: the prevalence of elevated blood pressure grew consistently with higher BMI percentiles. This trend appears to indicate a connection that is not solely dependent on regular physical development. This observation implies that obesity functions as an autonomous risk factor for hypertension, underscoring its independent role in contributing to high blood pressure.²⁰ Our results confirm the findings of Erlingsdottir *et al.*²¹ who showed an association between overweight/obesity and higher blood pressure. The mechanism whereby overweight/obesity may lead to higher blood pressure seems to be due to

enhanced adipocyte secretion of adipokines and pro-inflammatory cytokines which may disrupt normal physiological function leading to increased blood pressure.²¹

Age and gender

In the current study, Age was not associated with hypertension. Similar findings have been documented by Oyewole and Oritogun.²² but disagree with a Nigerian study that found age was associated with increases in blood pressure among adolescents, as well as the mid-adolescent age group (13-15 years).²³ The present study indicated that obese and overweight boys had a larger prevalence risk of high normal BP or hypertension than counterpart girls. The finding among Portuguese school adolescents aged 10 – 18 also showed a higher risk of being hypertensive among obese and overweight boys than girls.²⁴

In addition, the prevalence of hypertension in boys was greater than in girls, and this may be due to the role of testosterone in hypertension observed in males compared to non-menopausal women, which may explain the cause of hypertension in boys in this study.^{25,26}

The reason for girls having low blood pressure compared to boys may be attributed to estrogen and its protective effect on the cardiovascular system, because estrogen causes vasodilatation by modulating the function of vascular endothelial cells.^{27,28} The other probable cause of high blood pressure in adolescents aged (12-18) years could be a persistent hyperactive sympathetic nervous system even at rest which increases the smooth muscle tone of the vessels increasing resistance and hence persistently elevating pulse rate, cardiac output and hence elevating blood pressure.^{29,30}

Recommendations

Raising awareness among school students, particularly adolescents, regarding hypertension and its associated complications holds paramount importance. To achieve this objective, educational initiatives should be consistently organized by school authorities. These programs have the potential to inspire students to adopt healthier lifestyles and dietary habits. Additionally, such efforts can contribute to enhancing parents' understanding of hypertension, particularly when it is in its early stages.

Screening procedures should be implemented to detect asymptomatic hypertension in children, with a specific focus on mitigating influential factors such as obesity. This can be accomplished by offering guidance on balanced diets and regular physical activity, both within the home environment and at schools. Parents should also be actively encouraged to integrate these measures into their children's routines.

To this end, the Task Force on Blood Pressure Control in Children advocates for annual blood pressure measurements for all children aged 3 and above. These regular assessments hold significant value, as they can effectively identify hypertension in children. By collectively adhering to these practices, we can proactively address the issue of hypertension among the younger population.

Conclusions

This study showed that the prevalence of different classes of abnormal blood pressure was high in secondary school children in this region. The higher prevalence of hypertension among secondary school children was associated with being obese, using free fast in food, low level of education of parents. But being older and obesity predicted the prevalence of hypertension among secondary school children in this region.

References

1. Kearney PM, Whelton M, Reynolds K, et al. Global burden of hypertension: analysis of worldwide data. *Lancet* 2005;365:217-23.
2. Silverstein DM, Champoux E, Aviles DH, Vehaskari VMJPN. Treatment of primary and secondary hypertension in children. *Pediatr Nephrol* 2006;21:820-7.
3. Charan J, Buch N, Goyal JP, et al. Prevalence of hypertension in school going children of Surat city, Western India. *J Cardiovasc Dis Res* 2011;2:228-32.
4. Chobanian AV, Bakris GL, Black HR, et al. Seventh report of the joint national committee on prevention, detection, evaluation, and treatment of high blood pressure. *Hypertension* 2003;42:1206-52.

5. Alikhani S, Delavari A, Alaedini F, et al. A province-based surveillance system for the risk factors of non-communicable diseases: A prototype for integration of risk factor surveillance into primary healthcare systems of developing countries. *Public Health* 2009;123:358-64.
6. Subhi MD. Blood pressure profiles and hypertension in Iraqi primary school children. *Saudi Med J* 2006;27:482-6.
7. Grummer-Strawn LM, Reinold C, Krebs NF; Centers for Disease Control and Prevention (CDC). Use of World Health Organization and CDC growth charts for children aged 0-59 months in the United States. *MMWR Recomm Rep* 2010;59:1-15. Erratum in: *MMWR Recomm Rep* 2010;59:1184.
8. Goran MI, Ball GD, Cruz MLJTJoCE, Metabolism. Obesity and risk of type 2 diabetes and cardiovascular disease in children and adolescents. *J Clin Endocrinol Metab* 2003;88(4):1417-27.
9. Monge R, Beita OJJoah. Prevalence of coronary heart disease risk factors in Costa Rican adolescents. *J Adolesc Health* 2000;27:210-7.
10. Braveman P, Tarimo E. Screening in primary health care: setting priorities with limited resources: World Health Organization; 1994.
11. Raja T, Muthukumar T, Mohan AP. A cross sectional study on prevalence of hypertension and its associated risk factors among rural adults in Kanchipuram district, Tamil Nadu. *Int J Community Med Public Health* 2017;5:249-53.
12. Macedo ME, Lima MJ, Silva AO, et al. Prevalence, awareness, treatment and control of hypertension in Portugal: the PAP study. *J Hypertens* 2005;23:1661-6.
13. Jafar TH. Children, obesity, and high blood pressure: Asian populations at high risk. *Am J Hypert* 2009;22:6-7.
14. Falkner B. Hypertension in children and adolescents: epidemiology and natural history. *Pediatr Nephrol* 2010;25:1219-24.
15. Harrabi I, Belarbia A, Gaha R, et al. Epidemiology of hypertension among a population of school children in Sousse, Tunisia. *Can J Cardiol* 2006;22:212-6.

16. Kidy F, Rutebarika D, Lule SA, et al. Blood pressure in primary school children in Uganda: a cross-sectional survey. *BMC Public Health* 2014;14:1-9.
17. Nur N, Çetinkaya S, Yilmaz A, Ayvaz A, Bulut MO, Sümer HJJoh, population,, et al. Prevalence of hypertension among high school students in a middle Anatolian province of Turkey. *J Health Popul Nutr* 2008;26:88.
18. Sorof JM, Lai D, Turner J, et al. Overweight, ethnicity, and the prevalence of hypertension in school-aged children. *Pediatrics* 2004;113:475-82.
19. Karatzi K, Protogerou A, Rarra V, Stergiou GJ. Home and office blood pressure in children and adolescents: the role of obesity. The Arsakeion School Study. *J Hum Hypertens* 2009;23:512-20.
20. Meng L, Liang Y, Liu J, et al. Prevalence and risk factors of hypertension based on repeated measurements in Chinese children and adolescents. *Blood Press* 2013;22:59-64.
21. Erlingsdottir A, Indridason OS, Thorvaldsson O, Edvardsson VO. Blood pressure in children and target-organ damage later in life. *Pediatr Nephrol* 2010;25:323-8.
22. Oyewole O, Oritogun KS. Pre-hypertension and hypertension in adolescence: how much does it occur in a Nigerian community? *West Afr J Med* 2012;31:71-5.
23. Ujunwa FA, Ikefuna AN, Nwokocha AR, Chinawa JM. Hypertension and prehypertension among adolescents in secondary schools in Enugu, South East Nigeria. *Ital J Pediatr* 2013;39:1-6.
24. Rebelo D, Teixeira J, Marques-Vidal P, Oliveira JM. Obesity markers and blood pressure in a sample of Portuguese children and adolescents. *Eur J Cardiovasc Prev Rehabil* 2008;15:73-7.
25. Palmieri D, Perego P, Palombo DJA. Estrogen receptor activation protects against TNF- α -induced endothelial dysfunction. *Angiology* 2014;65:17-21.
26. Kolovou G, Giannakopoulou V, Vasiliadis Y, Bilianou H. Effects of estrogens on atherogenesis. *Curr Vasc Pharmacol* 2011;9:244-57.

27. Mendelsohn ME, Karas RH. The protective effects of estrogen on the cardiovascular system. *N Engl J Med* 1999;340:1801-11.
28. Alhalaiqa F, Abu-shbeeb I, Batiha A-M, Masa'Deh R, Amarneh B. The relation of demographic characteristics with fatigue levels among coronary heart disease patients: A Jordanian study. *Adv Stud Biol* 2015;7:301-22.
29. Grassi G, Mark A, Esler M. The sympathetic nervous system alterations in human hypertension. *Circ Res* 2015;116:976-90.
30. Kalil GZ, Haynes WGJHR. Sympathetic nervous system in obesity-related hypertension: mechanisms and clinical implications. *Hypertens Res* 2012;35:4-16.

Table 1. Prevalence of hypertension by age and gender among secondary school children.

General characteristeis	Statstics no (%)
Hypertension	
Abnormal blood pressure	85 (15.04)
Normotensive	480 (84.96)
Hypertension	
Normotensive	480 (84.96)
Elevated blood pressure	42 (7.43)
Hypertensive	33 (5.84)
Stage 1 hypertensive	4 (0.71)
Stage 2 hypertensive	6 (1.06)
Gender	
Male	242 (42.83)
Female	323 (57.17)
Age groups	
13	74 (13.10)
14	122 (21.59)
15	89 (15.75)
16	98 (17.35)
17	76 (13.45)
18	106 (18.76)
Residency	
Urban	57 (10.09)
Rural	508 (89.91)
Family size	
01-mar	79 (13.98)
04-giu	337 (59.65)
7 and above	149 (26.37)

Table 2. Prevalence of hypertension by age and gender among secondary school children.

Characteristics (n=565)	Blood pressure no (%)					p- value
	Normotensive	Elevated blood pressure	Hypertensive	Stage 1 hypertensive	Stage 2 hypertensive	
Hypertension	480 (84.96)	42 (7.43)	33 (5.84)	4 (0.71)	6 (1.06)	
Gender						0.0934
Male	205 (84.71)	18 (7.44)	11 (4.55)	4 (1.65)	4 (1.65)	
Female	275 (85.14)	24 (7.43)	22 (6.81)	0 (0.00)	2 (0.62)	
Age groups						0.2338
13	66 (89.19)	6 (8.11)	2 (2.70)	0 (0.00)	0 (0.00)	
14	104 (85.25)	7 (5.74)	11 (9.02)	0 (0.00)	0 (0.00)	
15	73 (82.02)	10 (11.24)	4 (4.49)	1 (1.12)	1 (1.12)	
16	86 (87.76)	5 (5.10)	5 (5.10)	2 (2.04)	0 (0.00)	
17	59 (77.63)	8 (10.53)	6 (7.89)	0 (0.00)	3 (3.95)	
18	92 (86.79)	6 (5.66)	5 (4.72)	1 (0.94)	2 (1.89)	
Residency						0.8967
Urban	433 (85.24)	37 (7.28)	29 (5.71)	4 (0.79)	5 (0.98)	
Rural	47 (82.46)	5 (8.77)	4 (7.02)	0 (0.00)	1 (1.75)	
Family size						0.7519
01-mar	65 (82.28)	7 (8.86)	6 (7.59)	1 (1.27)	0 (0.00)	
04-giu	282 (83.68)	27 (8.01)	21 (6.23)	2 (0.59)	5 (1.48)	
7 and above	133 (89.26)	8 (5.37)	6 (4.03)	1 (0.67)	1 (0.67)	
Pearson chi-squared test was performed for statistical analyses.						

Table 3. Association of school children blood pressure with socio-demographic characteristics of father.

Characteristics (n=565)	Blood pressure with father characteristics no (%)					p-value (two - sided)
	Elevated blood pressure	Hypertensive	Normotensive	Stage 1 hypertensive	Stage 2 hypertensive	
Education						0.0333
Illiterate	6 (8.82)	2 (2.94)	59 (86.76)	0 (0.00)	1 (1.47)	
Read and write	10 (6.90)	8 (5.52)	125 (86.21)	0 (0.00)	2 (1.38)	
Primary school	6 (4.58)	4 (3.05)	120 (91.60)	0 (0.00)	1 (0.76)	
Intermediately school	9 (12.50)	7 (9.72)	54 (75.00)	1 (1.39)	1 (1.39)	
Secondary	1 (3.57)	6 (21.43)	21 (75.00)	0 (0.00)	0 (0.00)	
Institute/college and above	10 (9.17)	6 (5.50)	89 (81.65)	3 (2.75)	1 (0.92)	
Overweight/obesity						0.0421
No	24 (5.76)	22 (5.28)	365 (87.53)	3 (0.72)	3 (0.72)	
Yes	18 (12.16)	11 (7.43)	115 (77.70)	1 (0.68)	3 (2.03)	
Hypertension						0.0345
No	36 (8.09)	23 (5.17)	381 (85.62)	1 (0.22)	4 (0.90)	
Yes	6 (5.00)	10 (8.33)	99 (82.50)	3 (2.50)	2 (1.67)	
Sleep apnea						0.0002
No	41 (7.45)	32 (5.82)	469 (85.27)	4 (0.73)	4 (0.73)	
Yes	1 (6.67)	1 (6.67)	11 (73.33)	0 (0.00)	2 (13.33)	
Blood pressure and mother characteristics no (%)						
Education						0.0291
Illiterate	13 (6.60)	13 (6.60)	168 (85.28)	1 (0.51)	2 (1.02)	
Read and write	7 (4.76)	6 (4.08)	133 (90.48)	0 (0.00)	1 (0.68)	
Primary school	11 (12.09)	5 (5.49)	74 (81.32)	1 (1.10)	0 (0.00)	

Intermediately school	2 (4.35)	1 (2.17)	42 (91.30)	0 (0.00)	1 (2.17)	0.0083
Secondary	0 (0.00)	1 (4.76)	19 (90.48)	0 (0.00)	1 (4.76)	
Institute/college	9 (15.79)	7 (12.28)	38 (66.67)	2 (3.51)	1 (1.75)	
Occupation						
Housewife	33 (6.61)	28 (5.61)	431 (86.37)	2 (0.40)	5 (1.00)	
Employee	9 (15.00)	5 (8.33)	43 (71.67)	2 (3.33)	1 (1.67)	
Pearson chi-squared test was performed for statistical analyses.						
Non significant factors were not presented in this table.						

Table 4. Association of hypertension with lifestyle related factors among school children.

Life style factors (n=565)	Blood pressure no (%)					p-value (two-sided)
	Normotensive	Elevated blood pressure	Hypertensive	Stage 1 hypertensive	Stage 2 hypertensive	
Exercise						0.3080
No	234 (83.27)	20 (7.12)	19 (6.76)	3 (1.07)	5 (1.78)	
Yes	246 (86.62)	22 (7.75)	14 (4.93)	1 (0.35)	1 (0.35)	
Smoking						0.1754
No	404 (85.05)	35 (7.37)	29 (6.11)	4 (0.84)	3 (0.63)	
Yes	76 (84.44)	7 (7.78)	4 (4.44)	0 (0.00)	3 (3.33)	
Sleep						0.8568
Short sleeper	18 (78.26)	2 (8.70)	2 (8.70)	0 (0.00)	1 (4.35)	
Normal sleeper	278 (84.76)	24 (7.32)	21 (6.40)	2 (0.61)	3 (0.91)	
Long sleeper	184 (85.98)	16 (7.48)	10 (4.67)	2 (0.93)	2 (0.93)	
Breakfast						0.2382
no	162 (88.04)	9 (4.89)	10 (5.43)	0 (0.00)	3 (1.63)	
yes	318 (83.46)	33 (8.66)	23 (6.04)	4 (1.05)	3 (0.79)	
Lunch						0.6383
no	77 (81.91)	10 (10.64)	6 (6.38)	0 (0.00)	1 (1.06)	
yes	403 (85.56)	32 (6.79)	27 (5.73)	4 (0.85)	5 (1.06)	

Dinner						0.4956
no	53 (89.83)	5 (8.47)	1 (1.69)	0 (0.00)	0 (0.00)	
yes	427 (84.39)	37 (7.31)	32 (6.32)	4 (0.79)	6 (1.19)	
Snack						0.5333
no	218 (82.89)	22 (8.37)	16 (6.08)	3 (1.14)	4 (1.52)	
yes	262 (86.75)	20 (6.62)	17 (5.63)	1 (0.33)	2 (0.66)	
Junk food						0.4753
no	41 (83.67)	6 (12.24)	2 (4.08)	0 (0.00)	0 (0.00)	
1-3 time a week	247 (84.30)	25 (8.53)	16 (5.46)	1 (0.34)	4 (1.37)	
> 3 times week	192 (86.10)	11 (4.93)	15 (6.73)	3 (1.35v)	2 (0.90)	
Type of foods						0.5178
fast food	40 (83.33)	2 (4.17)	4 (8.33)	1 (2.08)	1 (2.08)	
Homemade food	440 (85.11)	40 (7.74)	29 (5.61)	3 (0.58)	5 (0.97)	
Sugar sweeter beverage						0.8457
No	49 (84.48)	5 (8.62)	4 (6.90)	0 (0.00)	0 (0.00)	
Yes	431 (85.01)	37 (7.30)	29 (5.72)	4 (0.79)	6 (1.18)	
Salt consumption						0.6364
mild to moderate	439 (85.08)	39 (7.56)	29 (5.62)	3 (0.58)	6 (1.16)	
freely used by family	41 (83.67)	3 (6.12)	4 (8.16)	1 (2.04)	0 (0.00)	
Fat consumption						0.0375
mild moderate	397 (85.38)	33 (7.10)	29 (6.24)	1 (0.22)	5 (1.08)	
Freely used by family	83 (83.00)	9 (9.00)	4 (4.00)	3 (3.00)	1 (1.00)	
Past medical history						0.9255
No	475 (84.82)	42 (7.50)	33 (5.89)	4 (0.71)	6 (1.07)	
Yes	5 (100.00)	0 (0.00)	0 (0.00)	0 (0.00)	0 (0.00)	
Past surgical history						0.4252
No	413 (84.46)	38 (7.77)	30 (6.13)	4 (0.82)	4 (0.82)	
Yes	67 (88.16)	4 (5.26)	3 (3.95)	0 (0.00)	2 (2.63)	
BMI						<0.0001
Underweight	40 (97.56)	1 (2.44)	0 (0.00)	0 (0.00)	0 (0.00)	

Normal weight	342 (93.19)	15 (4.09)	9 (2.45)	0 (0.00)	1 (0.27)	
Overweight	57 (68.67)	12 (14.46)	11 (13.25)	1 (1.20)	2 (2.41)	
Obese	41 (55.41)	14 (18.92)	13 (17.57)	3 (4.05)	3 (4.05)	
Pearson chi-squared test was performed for statistical analyses.						

Table 5: Predictors of hypertension in school-aged children


















Predictors	Outcome: Blood pressure	P-value
BMI percentage		0.00000
Student Age		0.00051
Break fast		0.21217
Dinner		0.22074
Fat consumption		0.35812
Exercise		0.39232
Salt consumption		0.55886
Gender		0.64187
Type of foods		0.67822
Snack		0.71151
Family size category		0.76410
Junk food frequency		0.77417
Lunch		0.83531
Sugar sweeter beverage		0.83846
Smoking		0.86392
Residence		0.94736
Sleep duration category		0.99309
Nominal logistic regression was performed for statistical analyses.		

Table 5. Association of blood pressure with quality of life among secondary school children.

Life style factors (n=565)	Blood pressure no (%)					p-value (two-sided)
	Normotensive	Elevated blood pressure	Hypertensive	Stage 1 hypertensive	Stage 2 hypertensive	
QoL	23.85 (4.31)	23.57 (5.23)	24.00 (4.18)	28.25 (5.25)	26.17 (3.71)	0.2103

The comparisons were not statistically significant by gender as well.
ANOVA one-way was performed for statistical analysis.




Figure 1. Prevalence of hypertension among children with different BMI.




Figure 2. Comparisons of QoL among school children with blood pressure category.