



Italian Journal of Food Safety

<https://www.pagepressjournals.org/index.php/ijfs/index>

eISSN 2239-7132

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Italian Journal of Food Safety** is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Please cite this article as:

Lorusso P, Pandiscia A, Manfredi A, et al. **Evaluation of hepatitis E virus RNA persistence in experimentally contaminated cured pork liver sausages.** *Ital J Food Saf*
doi:10.4081/ijfs.2024.12286

Submitted: 15-01-2024

Accepted: 11-03-2024

© the Author(s), 2024
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.

All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Evaluation of hepatitis E virus RNA persistence in experimentally contaminated cured pork liver sausages

Patrizio Lorusso,¹ Annamaria Pandiscia,¹ Alessio Manfredi,¹
Giuseppina Marilia Tantillo,² Valentina Terio¹

¹Department of Veterinary Medicine, University of Bari; ²Department of Interdisciplinary Medicine, University of Bari, Italy

Correspondence: Patrizio Lorusso, Department of Veterinary Medicine, University of Bari, Provincial Road to Casamassima Km 3, 70010 Valenzano (Bari), Italy.

E-mail: patrizio.lorusso@uniba.it

Key words: HEV-3, food-borne viruses, liver sausage, virus persistence, cured meat product.

Contributions: PL, conceptualization, investigation, writing - original draft; AP, investigation, writing - review and editing; AM, investigation, writing - original draft; GMT, conceptualization, supervision; VT, conceptualization, writing - review and editing, supervision

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: not applicable.

Funding: none.

Availability of data and materials: data and materials are available from the corresponding author upon request.

Conference presentation: this paper was presented at the XXXII National Conference of the Italian Association of Veterinary Food Hygienists (AIVI), Maierato-Vibo Valentia, September 13-14-15, 2023.

Acknowledgments: the authors would like to thank Dr. Reimar Jhon, German Federal Institute for Risk Assessment, Department of Biological Safety, for kindly providing us with the persistently infected A549 cell lines and the cell culture adapted HEV genotype 3 strain 47832c.

Abstract

Hepatitis E is a disease sustained by RNA viruses, which have four different genotypes, all of which are responsible for acute forms of hepatitis. Genotypes 1 and 2 infect only humans, causing epidemics mainly transmitted by contaminated water, while genotypes 3 and 4 are zoonotic, and the infection is linked to the consumption of raw or undercooked meat or meat products. Hepatitis E virus (HEV) genotypes 3 and 4 have been detected in domestic Suidae, considered the asymptomatic reservoir of HEV, and in wild animals such as wild boar and deer. Despite scientific studies that have highlighted the presence of HEV in cured meat products, such as pork liver sausages, the viral persistence in the different production steps of curing has not been evaluated. Therefore, this study aimed to evaluate the persistence of HEV genotype 3 during the different curing and storage times of experimentally contaminated pork liver sausages using biomolecular methods. The sausages tested positive at all curing and storage times. This study confirms the potential risk attributed to pork liver sausages in HEV transmission. However, to guarantee an efficient risk assessment, future studies will be performed to correlate the presence of HEV RNA with infectious viral particles.

Introduction

Hepatitis E is an acute viral disease caused by the hepatitis E virus (HEV) that is classified in the family Hepeviridae. Strains affecting humans belong to the *Orthohepevirus A* species, recently re-named *Paslahepevirus balayani* from the *Paslahepevirus* genus, according to the new taxonomy classification by the International Committee on Taxonomy of Viruses (n.d.).

The characterization of the *P. balayani* genome showed eight different genotypes (HEV-1 to HEV-8) and only one serogroup (Smith *et al.*, 2020). HEV-1 and HEV-2 are linked only to human infections, and they have been associated with significant outbreaks and epidemics, particularly in developing countries (Pallerla *et al.*, 2020). Instead, HEV-3 and HEV-4 are associated with zoonotic infections that cause small outbreaks (Said *et al.*, 2009; Garbuglia *et al.*, 2013; Hakim *et al.*, 2017). These infections typically occur when consumers eat raw or undercooked food of animal origin, such as sausages, made with meat or liver from infected wild or domestic animals (boar, pigs, and deer) (Tei *et al.*, 2003; Yazaki *et al.*, 2003; Masuda *et al.*, 2005; Deest *et al.*, 2007; Colson *et al.*, 2010; Ricci *et al.*, 2017). HEV-4 is spread in East Asia, while HEV-3 was detected worldwide and is recognized as an emerging pathogen in Western countries, including Europe, where, over the past decade, there have been more than 21,000 cases with 28 deaths (Ricci *et al.*, 2017). In particular, there is an increasing trend of cases of HEV infection in Italy, mainly in central-northern Italy (Istituto Superiore di Sanità, n.d.). In Italy, the HEV seroprevalence detected in blood donors showed low to moderate rates (from 1.3% up to 27.20%). High levels in some regions and/or provinces were mainly attributable to eating habits (Spada *et al.*, 2022).

In Italy, the consumption of raw or undercooked pig/wild boar meat and liver sausages represents one of the main risk factors linked to human HEV infection (Garbuglia *et al.*, 2021). In fact, the consumption of pork meat and its products increases every year, reaching 18.8 kg. pro capite (Lenti, 2022). The incubation periods (30-60 days) of this virus do not allow for a correlate of the infection with the food vehicle responsible for food-borne illness (Garbuglia *et al.*, 2021). Moreover, another critical point is that most of the suspected food was no longer available for the HEV investigation, at the onset of symptoms. So, to better understand the role of pork products in the transmission of HEV, this study aims to investigate the persistence of HEV RNA in artificially contaminated pork liver sausages cured and stored at different time intervals.

Materials and Methods

Virus stock preparation

The cell culture-adapted HEV genotype 3 strain 47832c (GenBank acc. no. KC618403), originally isolated from a serum sample of a chronically HEV-infected patient (Johne *et al.*, 2014), was used in all experiments. A cell line of A549 cells persistently infected with this virus strain (Johne *et al.*, 2014; Johne *et al.*, 2016) was used for the production of the virus. The persistently HEV-infected A549 cell

line was grown in Dulbecco's modified eagle medium with 10% fetal bovine serum (FBS, 10270106, Gibco, Waltham, MA, USA) and 100 µg/mL gentamicin (15750060, Gibco, Waltham, MA, USA) at 37°C and 5% CO₂ in a humidified incubator. After 7 days, some of the flasks were subjected to three cycles of freezing (-20°C) and thawing [room temperature (RT)] to release non-enveloped virus particles from inside the cells into the supernatant. The resulting supernatant was stored at -20°C. The viral load was evaluated with real-time polymerase chain reaction (PCR).

Sausage preparation

A total of 11 liver sausages were prepared in collaboration with a local sausage company. The liver sausages were prepared according to the following recipe: 60% pork, 40% pork liver, and 2.5% sea salt. All pork meat and liver were previously tested with biomolecular methods to assess the absence of HEV RNA. Fresh sausages were uniformly infected with a viral suspension containing 1000 GC/g by injection along the entire sausage. One sausage sample was used as the start point control (without curing and storing, T₀). Subsequently, all sausages were cured at a temperature between 20°C and 25°C. Six sausage samples (Tc 10, Tc 15, Tc 20, Tc 25, Tc 30, Tc 40) were analyzed after 10, 15, 20, 25, 30, and 40 curing days. Moreover, four sausage samples (Ts 20, Ts 30, Ts 40, Ts 45) cured for 40 days, were stored at 4 °C for 20, 30, 40, and 45 days.

Sample preparation and nucleic acid extraction

Virus extraction was performed following the method described by Di Pasquale *et al.* (2019) with appropriate modifications. Briefly, 1 g of liver sausage was shredded with a sterile blade and added to a tube containing 3.5 mL of TRIZOL Reagent (Life Technologies, Monza, Italy) and 1 sterile tungsten carbide bead (3 mm diameter). Samples were homogenized for 5 minutes at 25 hz/s in Tissue Lyser II (Qiagen, GmbH, Hilden, Germany). After mechanical disruption of the livers, samples were incubated at RT for 15 minutes and centrifugated at 8000× g for 20 minutes at 4°C. Then the supernatant was added to 0.7 mL of chloroform, vortexed for 15 seconds, and incubated for 15 minutes at RT. The aqueous phase obtained by centrifugation at 8000× g for 15 minutes at 4°C, was recovered and stored at -80°C until use.

Touchdown reverse transcription-polymerase chain reaction and touchdown hemi-nested polymerase chain reaction

The touchdown reverse transcription-PCR (TD/RT-PCR) and assay-touchdown hemi-nested PCR (TD/hn-PCR) were carried out as described in detail by Lorusso *et al.* (2022). Briefly, the RNA extract was retrotranscribed by TD/RT-PCR and subsequently amplified with TD/hn-PCR to increase sensitivity using primers reported by Drexler *et al.* (2012), amplifying a fragment of 346 bp belonging to RdRp of ORF1.

Detection of amplified products

TD/hn-PCR-amplified products were displayed by electrophoresis on 1.5% (w/v) agarose NA (Pharmacia, Uppsala, Sweden) gel in 1X TBE buffer containing 0.089 M Tris, 0.089 M boric acid, 0.002 M EDTA, pH 8.0 (USB, Cleveland, OH, USA), and stained with Green Gel Safe 10000X Nucleic Acid Stain (5 l/100 ml) (Fisher Molecular Biology, USA). The Gene Ruler™ 100 bp DNA Ladder molecular weight marker (MBI Fermentas, Vilnius, Lithuania) was used. Image acquisition was performed with Gel Doc™ EZ imager Bio-rad (Bio-Rad Laboratories, Inc. Hercules, California, USA).

Results

All sausage samples analyzed at different curing and storage times tested positive for HEV RNA, as described in Table 1.

Discussion

In our study, all pork liver sausages experimentally contaminated with HEV viral particles tested positive for HEV RNA at all curing and storage times (Table 1). Despite our study being focused on experimentally contaminated pork liver sausages, the results obtained have been enforced by previous studies on liver sausages marketed in Europe that reported a prevalence of HEV RNA from 4.3% to 70% (Giannini *et al.*, 2018; Boxman *et al.*, 2019; Montone *et al.*, 2019). In addition, these studies suggest that the curing process potentially preserves the HEV RNA (Di Bartolo *et al.*, 2015).

Although curing is a technique useful in increasing the shelf life of food by reducing its pathogenic and altering microbial load, it has been demonstrated that it is not able to reduce the viral load (Sankaranarayanan *et al.*, 2019; Colavita, 2023). Especially, the antimicrobial action of curing is mainly due to the reduction of activity in water and the lowering of pH (Colavita, 2023). Previous studies reported that food-borne viruses are characterized by high resistance to extreme pH, drying and thermal stress, and additives normally used in sausages since they are without envelope (Straube *et al.*, 2011; Johne *et al.*, 2016; Bosch *et al.*, 2018; Wolff *et al.*, 2020; Wolff *et al.*, 2022). To date it is well known that cooking contaminated pork or wild boar meat and their meat products, at 71°C for no less than 20 minutes, is useful in reducing the risk of HEV infection (Barnaud *et al.*, 2012; Ricci *et al.*, 2017).

Our study detected a high persistence of HEV RNA in pork liver sausages. It could be due to the presence of fat in their composition, which could act as protection against degrading enzymes of viral particles, such as proteases and RNAsi. This hypothesis, however, requires further investigations that allow us to understand the role of fat in the potential protection of HEV viral particles and how/if it could interact with them.

Moreover, we used reagents dissolving the fat portion to improve the extraction efficiency since it could prevent the extraction step of viral particles, producing false negative results (Sair *et al.*, 2002; Martínez-Martínez *et al.*, 2011; Schrader *et al.*, 2012; Szabo *et al.*, 2015).

In the present study, we evaluated only the occurrence of viral RNA without determining the presence of infectious viral particles in the liver pork sausages analyzed (Kamar *et al.*, 2014). Nevertheless, when foods have a high GC/g of viral particles, we can assume that there are also infectious particles (Giannini *et al.*, 2018). To date, the infectious dose of HEV is not known, but the severity of the disease depends on the presence of infectious viral particles and the host immunity (Tsarev *et al.*, 1994; Kamar *et al.*, 2014), particularly for fragile individuals such as immunocompromised persons, pregnant women, and persons with liver disease (Giannini *et al.*, 2018).

Finally, to guarantee an efficient risk assessment, a future study will be performed to correlate the presence of HEV RNA with infectious viral particles.

Conclusions

Our study shows that curing is not an effective method to eliminate the presence of HEV RNA in pork liver sausages. Therefore, to prevent potential epidemic outbreaks of HEV due to the intake of meat and/or liver products consumed raw or undercooked, several approaches should be followed: i) educating consumers through information campaigns regarding the consumption of pork and wild boar meat and sausages cooked; ii) providing systematic sampling to evaluate the presence of HEV RNA in meat products to ensure the consumers' health.

Reference

- Barnaud E, Rogée S, Garry P, Rose N, Pavio N, 2012. Thermal inactivation of infectious hepatitis E virus in experimentally contaminated food. *Appl Environ Microbiol* 78:5153-9.
- Bosch A, Gkogka E, Le Guyader FS, Loisy-Hamon F, Lee A, van Lieshout L, Marthi B, Myrmel M, Sansom A, Schultz AC, Winkler A, Zuber S, Phister T, 2018. Foodborne viruses: detection, risk assessment, and control options in food processing. *Int J Food Microbiol* 285:110-28.

- Boxman ILA, Jansen CCC, Hägele G, Zwartkruis-Nahuis A, Tijsma ASL, Vennema H, 2019. Monitoring of pork liver and meat products on the Dutch market for the presence of HEV RNA. *Int J Food Microbiol* 296:58-64.
- Colavita G, 2023. Igiene e tecnologie degli alimenti. Point Veterinaire Italie, Milan, Italia.
- Colson P, Borentain P, Queyriaux B, Kaba M, Moal V, Gallian P, Heyries L, Raoult D, Gerolami R, 2010. Pig liver sausage as a source of hepatitis E virus transmission to humans. *J Infect Dis* 202: 825-34.
- Deest G, Zehner L, Nicand E, Gaudy-Graffin C, Goudeau A, Bacq Y, 2007. Autochthonous hepatitis E in France and consumption of raw pig meat. *Gastroenterol Clin Biol* 31:1095-7.
- Di Bartolo I, Angeloni G, Ponterio E, Ostanello F, Ruggeri FM, 2015. Detection of hepatitis E virus in pork liver sausages. *Int J Food Microbiol* 193:29-33.
- Di Pasquale S, De Santis P, La Rosa G, Di Domenico K, Iaconelli M, Micarelli G, Martini E, Bilei S, De Medici D, Suffredini E, 2019. Quantification and genetic diversity of Hepatitis E virus in wild boar (*Sus scrofa*) hunted for domestic consumption in Central Italy. *Food Microbiol* 82: 194-201.
- Drexler JF, Seelen A, Corman VM, Fumie Tateno A, Cottontail V, Melim Zerbinati R, Gloza-Rausch F, Klose SM, Adu-Sarkodie Y, Oppong SK, Kalko EKV, Osterman A, Rasche A, Adam A, Müller MA, Ulrich RG, Leroy EM, Lukashev AN, Drosten C, 2012. Bats worldwide carry hepatitis E virus-related viruses that form a putative novel genus within the family Hepeviridae. *J Virol* 86.
- Garbuglia AR, Bruni R, Villano U, Vairo F, Lapa D, Madonna E, Picchi G, Binda B, Mariani R, De Paulis F, D'amato S, Grimaldi A, Scognamiglio P, Capobianchi MR, Ciccaglione AR, 2021. Hepatitis E outbreak in the central part of Italy sustained by multiple HEV genotype 3 strains, June-December 2019. *Viruses* 13:1159.
- Garbuglia AR, Scognamiglio P, Petrosillo N, Mastroianni CM, Sordillo P, Gentile D, La Scala P, Girardi E, Capobianchi MR, 2013. Hepatitis E virus genotype 4 outbreak, Italy, 2011. *Emerg Infect Dis* 19:110-4.
- Giannini P, Jermini M, Leggeri L, Nüesch-Inderbinen M, Stephan R, 2018. Detection of hepatitis E virus RNA in raw cured sausages and raw cured sausages containing pig liver at retail stores in Switzerland. *J Food Prot* 81:43-5.
- Hakim MS, Wang W, Bramer WM, Geng J, Huang F, de Man RA, Peppelenbosch MP, Pan Q, 2017. The global burden of hepatitis E outbreaks: a systematic review. *Liver Int* 37:19-31.
- International Committee on Taxonomy of Viruses, n.d. Subfamily: Orthohepevirinae Genus: Paslahepevirus. Available from: <https://ictv.global/report/chapter/hepeviridae/hepeviridae/orthohepevirinae/paslahepevirus>. Accessed on: 27/12/2023.
- Istituto Superiore di Sanità, n.d. Epatite virale. Available from: <https://www.epicentro.iss.it/epatite/dati-seieva>. Accessed on: 27/12/23. [Material in Italian].
- Johne R, Reetz J, Ulrich RG, Machnowska P, Sachsenröder J, Nickel P, Hofmann J, 2014. An ORF1-rearranged hepatitis E virus derived from a chronically infected patient efficiently replicates in cell culture. *J Viral Hepatitis* 21:447-56.
- Johne R, Trojnar E, Filter M, Hofmann J, 2016. Thermal stability of hepatitis E virus as estimated by a cell culture method. *Appl Environ Microbiol* 82:4225-31.
- Kamar N, Dalton HR, Abravanel F, Izopet J, 2014. Hepatitis E virus infection. *Clin Microbiol Rev* 27:116-38.
- Lenti R, 2022. Rapporto Annuale 2022. Available from: <https://www.assica.it/news/#rapporto-annuale>.
- Lorusso P, Bonerba E, Pandiscia A, Mottola A, Di Pinto A, Piredda R, Terio V, 2022. Occurrence of hepatitis E virus (HEV) in Calabrian wild boars. *Int J Food Microbiology* 371:109671.
- Martínez-Martínez M, Díez-Valcarce M, Cook N, Hernández M, Rodríguez-Lázaro D, 2011. Evaluation of extraction methods for efficient detection of enteric viruses in pork meat products. *Food Anal Method* 4:13-22.

- Masuda JI, Yano K, Tamada Y, Takii Y, Ito M, Omagari K, Kohno S, 2005. Acute hepatitis E of a man who consumed wild boar meat prior to the onset of illness in Nagasaki, Japan. *Hepatology* 31:178-83.
- Montone AMI, De Sabato L, Suffredini E, Alise M, Zaccherini A, Volzone P, Di Maro O, Neola B, Capuano F, Di Bartolo I, 2019. Occurrence of HEV-RNA in Italian regional pork and wild boar food products. *Food Environ Virol* 11:420-6.
- Pallerla SR, Harms D, Johne R, Todt D, Steinmann E, Schemmerer M, Wenzel JJ, Hofmann J, Kuo Shih JW, Wedemeyer H, Bock CT, Velavan TP, 2020. Hepatitis E virus infection: circulation, molecular epidemiology, and impact on global health. *Pathogens* 9:856.
- Ricci A, Allende A, Bolton D, Chemaly M, Davies R, Fernandez Escamez PS, Herman L, Koutsoumanis K, Lindqvist R, Nørnung B, Robertson L, Ru G, Sanaa M, Simmons M, Skandamis P, Snary E, Speybroeck N, Ter Kuile B, Threlfall J, Wahlström H, Di Bartolo I, Johne R, Pavio N, Rutjes S, van der Poel W, Vasickova P, Hempen M, Messens W, Rizzi V, Latronico F, Girones R, 2017. Public health risks associated with hepatitis E virus (HEV) as a food-borne pathogen. *EFSA J* 15:e04664.
- Said B, Ijaz S, Kafatos G, Booth L, Thomas HL, Walsh A, Ramsay M, Morgan D, 2009. Hepatitis E outbreak on cruise ship. *Emerg Infect Dis* 15:1738-44.
- Sair AI, D'Souza DH, Moe CL, Jaykus LA, 2002. Improved detection of human enteric viruses in foods by RT-PCR. *J Virol Methods* 100:57-69.
- Sankaranarayanan A, Amaesan N, Dhanasekaran D, 2019. Fermented food products. CRC Press, Boca Raton, FL, USA.
- Schrader C, Schielke A, Ellerbroek L, Johne R, 2012. PCR inhibitors - occurrence, properties and removal. *J Appl Microbiol* 113:1014-26.
- Smith DB, Izopet J, Nicot F, Simmonds P, Jameel S, Meng XJ, Norder H, Okamoto H, van der Poel WHM, Reuter G, Purdy MA, 2020. Update: proposed reference sequences for subtypes of hepatitis E virus (species Orthohepevirus A). *J Gen Virol* 101:692-8.
- Spada E, Simeoni M, Martina A, Pati I, Villano U, Adriani D, D'Angiò A, Tritarelli E, Taffon S, Bellino S, Boros S, Urciuoli R, Masiello F, Marano G, Bruni R, Pezzotti P, Ciccaglione AR, Pupella S, De Angelis V, Pisani G, 2022. Prevalence and risk factors for hepatitis E virus infection in blood donors: a nationwide survey in Italy, 2017 to 2019. *Euro Surveill* 27:2100516.
- Straube J, Albert T, Manteufel J, Heinze J, Fehlhaber K, Truyen U, 2011. In vitro influence of D/L-lactic acid, sodium chloride and sodium nitrite on the infectivity of feline calicivirus and of ECHO virus as potential surrogates for foodborne viruses. *Int J Food Microbiol* 151:93-7.
- Szabo K, Trojnar E, Anheyer-Behmenburg H, Binder A, Schotte U, Ellerbroek L, Klein G, Johne R, 2015. Detection of hepatitis E virus RNA in raw sausages and liver sausages from retail in Germany using an optimized method. *Int J Food Microbiol* 215:149-56.
- Tei S, Kitajima N, Takahashi K, Mishihiro S, 2003. Zoonotic transmission of hepatitis E virus from deer to human beings. *Lancet* 362:371-3.
- Tsarev SA, Tsareva TS, Emerson SU, Govindarajan S, Shapiro M, Gerin JL, Purcell RH, 1994. Successful passive and active immunization of cynomolgus monkeys against hepatitis E. *Proc Natl Acad Sci U S A* 91:10198-202.
- Wolff A, Günther T, Albert T, Schilling-Loeffler K, Gadicherla AK, Johne R, 2020. Stability of hepatitis E virus at different pH values. *Int J Food Microbiol* 325:108625.
- Wolff A, Günther T, Johne R, 2022. Stability of hepatitis E virus after drying on different surfaces. *Food Environ Virol* 14:138-48.
- Yazaki Y, Mizuo H, Takahashi M, Nishizawa T, Sasaki N, Gotanda Y, Okamoto H, 2003. Sporadic acute or fulminant hepatitis E in Hokkaido, Japan, may be food-borne, as suggested by the presence of hepatitis E virus in pig liver as food. *J Gen Virol* 84:2351-7.

Table 1. Results of hepatitis E virus RNA analysis.

		Curing time						Storage time			
Day	T0	Tc10	Tc 15	Tc 20	Tc 25	Tc 30	Tc 40	Ts 20	Ts 30	Ts 40	Ts 45
HEV RNA test	+	+	+	+	+	+	+	+	+	+	+

Tc, curing time; Ts, storage time.