



eISSN 2239-7132

Italian Journal of Food Safety

<https://www.pagepressjournals.org/index.php/ijfs/index>

Publisher's Disclaimer. E-publishing ahead of print is increasingly important for the rapid dissemination of science. The Early Access service lets users access peer-reviewed articles well before print/regular issue publication, significantly reducing the time it takes for critical findings to reach the research community.

These articles are searchable and citable by their DOI (Digital Object Identifier).

The **Italian Journal of Food Safety** is, therefore, E-publishing PDF files of an early version of manuscripts that have undergone a regular peer review and have been accepted for publication, but have not been through the copyediting, typesetting, pagination, and proofreading processes, which may lead to differences between this version and the final one.

The final version of the manuscript will then appear in a regular issue of the journal.

The E-publishing of this PDF file has been approved by the authors.

Ital J Food Saf 2026 [Online ahead of print]

Please cite this article as:

Rampazzo G, Depau G, Pagliuca G, et al. **Cumulative exposure to per- and polyfluoroalkyl substances from eggs and egg products in Europe: application of relative potency factors.** *Ital J Food Saf* doi:10.4081/ijfs.2026.14673

Submitted: 25-11-2025

Accepted: 02-02-2026

 © the Author(s), 2026
Licensee PAGEPress, Italy

Note: The publisher is not responsible for the content or functionality of any supporting information supplied by the authors. Any queries should be directed to the corresponding author for the article.
All claims expressed in this article are solely those of the authors and do not necessarily represent those of their affiliated organizations, or those of the publisher, the editors and the reviewers. Any product that may be evaluated in this article or claim that may be made by its manufacturer is not guaranteed or endorsed by the publisher.

Cumulative exposure to per- and polyfluoroalkyl substances from eggs and egg products in Europe: application of relative potency factors

Giulia Rampazzo,¹ Giacomo Depau,¹ Giampiero Pagliuca,¹ Elisa Zironi,¹
Maria Federica Marchesi,¹ Francesco Arioli,² Teresa Gazzotti¹

¹Department of Veterinary Medical Sciences, Alma Mater Studiorum University of Bologna;
²Department of Veterinary Medicine and Animal Sciences, University of Milan, Lodi, Italy

Correspondence: Giulia Rampazzo, Department of Veterinary Medical Sciences, Alma Mater Studiorum University of Bologna, Via Tolara di Sopra 50, Ozzano dell'Emilia, 40064 Bologna, Italy.
E-mail: giulia.rampazzo4@unibo.it

Key words: PFAS, risk assessment, RPF, egg products, exposure assessment.

Contributions: all authors contributed equally.

Conflict of interest: the authors declare no potential conflict of interest.

Ethics approval and consent to participate: not applicable.

Availability of data and materials: data and materials are available from the corresponding author upon request.

Conference presentation: this paper was presented at the XXXIV National Conference of the Italian Association of Veterinary Food Hygienists (AIVI), Parma, September 10-11-12, 2025.

Abstract

Per- and polyfluoroalkyl substances (PFAS) are highly persistent contaminants frequently detected in foods of animal origin, raising concern for human health. Eggs and egg products are recognized by the European Food Safety Authority (EFSA) as one of the major dietary sources of PFAS exposure in Europe. This study applied the relative potency factors (RPFs), based on hepatic toxicity, to estimate cumulative dietary exposure to PFAS from eggs across different population groups. The 95th percentile occurrence data were combined with food consumption patterns to express exposure in perfluorooctanoic acid equivalents, providing a biologically relevant metric. Perfluorooctanesulfonic acid emerged as the dominant contributor to cumulative exposure. Toddlers exhibited the highest estimated weekly intake, reaching 1.63 ng/kg bw per week, while adolescents, adults, and elderly populations showed progressively lower values. These findings highlight children as a particularly vulnerable group due to higher intake per body weight. Consistent with previous studies, commercial eggs generally contained PFAS only at trace levels, whereas home-produced eggs often showed elevated contamination, in some cases exceeding the European Union maximum levels and EFSA's tolerable weekly intake. Overall, this work demonstrates the potential of the RPF approach for mixture-based risk assessment, underscores the need for refined exposure evaluation, and reinforces the importance of continuous monitoring of PFAS in eggs.

Introduction

Per- and polyfluoroalkyl substances (PFAS) are a large class of synthetic chemicals characterized by high stability and persistence, leading to widespread environmental and food contamination. Human exposure occurs mainly through diet, particularly animal-derived foods, which substantially contribute to internal body burdens (Domingo and Nadal, 2017; EFSA, 2020a; Genualdi *et al.*, 2022; Piva *et al.*, 2023; Nobile *et al.*, 2024; Depau *et al.*, 2025; Hossini *et al.*, 2025). In 2020, the European Food Safety Authority (EFSA) established a group tolerable weekly intake (TWI) of 4.4 ng/kg body weight (bw) per week for perfluorooctanoic acid (PFOA), perfluorononanoic acid (PFNA), perfluorooctane sulfonic acid (PFOS), and perfluorohexane sulfonic acid (PFHxS), based on immunotoxicity in children (EFSA, 2020a). These four PFAS were included due to their shared toxicokinetic and toxicological profiles, but they represent only a small subset of the thousands of PFAS identified (OECD, 2021). Some, such as PFOA and PFOS, have also been associated with hepatotoxicity, metabolic effects, developmental toxicity, and, more recently, carcinogenicity (Zahm *et al.*, 2024). Significant data gaps persist for less-studied PFAS, particularly emerging compounds, thereby limiting the robustness of consumer risk assessment. To improve risk assessment and reduce reliance on animal testing, European agencies (European Chemicals Agency, and EFSA) promote grouping and read-across strategies (European Chemicals Agency, 2016; EFSA, 2025). In mixture risk assessment, EFSA often applies the relative potency factor (RPF) approach, which expresses the potency of individual substances relative to an index compound, allowing cumulative exposure estimates in equivalent units (EFSA, 2012a; EFSA, 2020b). Bil *et al.* (2021) developed specific RPF for 23 PFAS using liver toxicity as the endpoint, enabling risk characterization in PFOA equivalents (PEQ) (Zeilmaker *et al.*, 2018; Bil *et al.*, 2021). Recent regulations, including the EU Drinking Water Directive (2020/2184) (European Parliament and Council of the European Union, 2020) and the Directive proposal on the protection of groundwater against pollution and deterioration (COM/2022/540) (European Commission, 2022), have already adopted sum parameters, RPF, and the PEQ concept, highlighting the growing regulatory emphasis on mixture assessment.

EFSA identifies eggs and egg products as one of the major sources of human PFAS dietary exposure (EFSA, 2020a). Several studies have reported the presence of PFAS in eggs, highlighting substantial differences between eggs produced in industrial systems (Gazzotti *et al.*, 2021; Chiumiento *et al.*, 2023; Nobile *et al.*, 2023) and those from backyard hens, with the latter often showing higher concentrations (Gazzotti *et al.*, 2021; Arvaniti *et al.*, 2025; Biesterbos and den Braver, 2025). Considering the increasing importance of evaluating as many PFAS as possible in food, as well as their potential implications for consumer health, the present study applies the RPF approach to

estimate the cumulative dietary exposure of European consumers to PFAS from eggs and egg-derived products, focusing on hepatic toxicity as the critical endpoint. By integrating occurrence and consumption data, and RPF, cumulative exposure is expressed in PEQs, providing a potentially more refined metric for exposure assessment. However, the RPF protocol proposed by Bil *et al.* (2021) has not yet received formal recognition or validation at the European level. As a result, some uncertainty persists, including the possibility of exposure overestimation (Rampazzo *et al.*, 2025). Further validation is required before this approach can be routinely applied in regulatory contexts.

Methods and Materials

Per- and polyfluoroalkyl substance occurrence, consumption, and consumer data

Occurrence data on PFAS in eggs and egg products were obtained from the EFSA 2020 scientific opinion on the risk to human health from PFAS in food and reported in Table 1(Annex A) (EFSA, 2020a). The 95th percentile lower bound occurrence values were used, in line with the realistic exposure assessment approach recommended by the EFSA CONTAM Panel (EFSA, 2020a). Consumption data were retrieved from the EFSA Comprehensive European Food Consumption Database. Average European consumption values were calculated as weighted means across Member State surveys for four age groups: toddlers, adolescents, adults, and the elderly and standard body weights (BW) for each group were applied as recommended by EFSA (Table 2) (EFSA, 2012b).

Perfluorooctanoic acid equivalents exposure evaluation

Occurrence data, combined with mean consumptions and bw, were converted to PEQ using RPFs (Table 3). Exposure was estimated as PEQ using minimum and maximum RPF values.

The estimated weekly intake (EWI) of each PFAS was calculated (1) for each consumer group and expressed in PEQ [Eq. 1]:

$$PEQ - EWI_{PFASi} = \frac{Concentration_i \times RPF_i(min/max) \times Mean Consumption}{BW} \times 7 \quad [Eq. 1]$$

Concentration_i = concentration of PFAS_i in eggs and egg products (ng/kg); RPF_i = RPF of PFAS_i; Mean Consumption = average daily consumption of eggs and egg products (kg/day); BW = standard body weight for the consumer group (kg); 7 = conversion from daily to weekly intake.

The cumulative PEQ weekly intake was then calculated [Eq. 2] for each consumer group:

$$Cumulative PEQ - EWI = \sum_{j=1}^n EWI_{PFASj} \quad [Eq. 2]$$

Results and Discussion

Table 4 shows the PEQ-EWI across different consumer groups. The use of the 95th percentile of occurrence data allows a focused assessment of the contribution of PFOA and PFOS to total PFAS exposure, which is particularly relevant given that the used data on PFAS occurrence in eggs and egg products are not normally distributed. In such distributions, a few highly contaminated samples can disproportionately elevate the mean. In contrast, the 95th percentile provides a more conservative and robust estimate of exposure, minimizing the influence of outliers. Across all age groups, PFOS remains the dominant contributor to exposure. Toddlers exhibit the highest cumulative PEQ-EWI, reaching 1.63 ng/kg bw per week, while adolescents, adults, and the elderly show progressively lower cumulative exposures, corresponding to 0.69, 0.53, and 0.50 ng/kg bw per week, respectively. The application of the RPF approach further strengthens these findings by highlighting the potentially greater contribution of PFOS to hepatic toxicity compared to PFOA, providing a more health-relevant perspective than simple concentration-based estimates.

Overall, this analysis underscores that children may represent a particularly vulnerable population due to their higher intake per body weight and the cumulative nature of PFAS. Even at low levels of individual compounds (considering the 4.4 ng/kg bw per week TWI value), their combined effects

may still pose a relevant risk, particularly given the recent IARC classifications of PFOA as carcinogenic and PFOS as possibly carcinogenic (Zahm *et al.*, 2024). These results are consistent with EFSA's assessment, which identifies eggs and egg products as one of the major sources of human PFAS dietary exposure (EFSA, 2020a). While studies on commercial eggs (Chiumiento *et al.*, 2023; Nobile *et al.*, 2023) detected only a few long-chain PFAS at trace levels, home-produced eggs (HPE) often show markedly higher contamination. Research in Italy (Gazzotti *et al.*, 2021), Belgium (Lasters *et al.*, 2022), and Greece (Arvaniti *et al.*, 2025) detected PFOS, PFOA, PFHxS, and PFBA at elevated concentrations, in some cases exceeding European maximum levels set for eggs and EFSA TWI. Backyard chickens are more exposed to contaminated soil, foraging, and feed, making HPE a significant contributor to PFAS dietary intake. Dutch risk assessments (Biesterbos and den Braver, 2024) indicate that PFAS exposure from HPE can be 20-80 times higher than from commercial eggs, further underscoring the potential health risks—particularly for children, who are more vulnerable due to higher intake per body weight. Moreover, expanding the RPF approach to include additional PFAS and critical endpoints (*e.g.*, immunotoxicity) would support the derivation of reference values for data-poor compounds. Integrating predictive methods within the RPF framework could enable more comprehensive and science-based mixture risk assessments, better reflecting real dietary exposure scenarios. Overall, these findings reinforce the importance of continuous monitoring of PFAS in food, the systematic application of mixture-based assessment approaches like RPF, and proactive management strategies aimed at safeguarding vulnerable populations—particularly children—against the potential cumulative and hepatic effects of PFAS in the diet.

Conclusions

In conclusion, this study confirms that eggs and egg products are an important source of dietary PFAS exposure and demonstrates the scientific potential of the RPF approach in dietary PFAS risk assessment. By combining compound-specific potency, occurrence data, and consumption patterns, the RPF framework allows cumulative exposure to be expressed in PEQ, potentially providing a more accurate and health-protective estimate than the traditional concentration-based method. This methodological distinction, however, reflects the proof-of-concept nature of the present study and underscores the need for further validation of the RPF approach before it can be routinely applied in risk assessment. In this context, PFOS may be identified as the dominant contributor to hepatic toxicity, with toddlers as the most vulnerable group due to higher PFAS intake per body weight. Beyond its role in exposure estimation, the RPF methodology offers a versatile and robust tool for mixture-based risk assessment and risk management, enabling the prioritization of compounds, identification of high-risk food sources, and the design of targeted mitigation strategies. Its implementation aligns with evolving European regulatory frameworks, which increasingly recognize the need to assess chemical mixtures rather than single substances, reflecting a shift toward more comprehensive and protective approaches in food safety.

References

- Arvaniti OS, Gerokonstantis DT, Bouzoukas C, Aloupi M, Gkotsis G, Konomi A, Mastrotheodoraki A, Iliopoulou A, Kostakis M, Dasenaki M, Thomaidis NS, Stasinakis AS, 2025. Occurrence of per- and polyfluoroalkyl substances, pesticides, pharmaceuticals, and heavy metals in Greek backyard chicken eggs and estimation of the consumption risk. *Sci Total Environ* 998:180253.
- Biesterbos JWH, den Braver MW, 2025. Risk assessment of PFAS in home-produced chicken eggs in the Netherlands. *FRAE* 3:0064E.
- Bil W, Zeilmaker M, Fragki S, Lijzen J, Verbruggen E, Bokkers B, 2021. Risk Assessment of per- and polyfluoroalkyl substance mixtures: a relative potency factor approach. *Environ Toxicol Chem* 40:859-70.
- Chiumiento F, Bellocchi M, Ceci R, D'Antonio S, De Benedictis A, Leva M, Pirito L, Rosato R, Scarpone R, Scortichini G, Tammaro G, Diletti G, 2023. A new method for determining

- PFASs by UHPLC-HRMS (Q-Orbitrap): application to PFAS analysis of organic and conventional eggs sold in Italy. *Food Chem* 401:134135.
- Depau G, Zampiga M, Rampazzo G, Zironi E, Sirri F, Pagliuca G, Gazzotti T, 2025. Preliminary results from monitoring perfluoroalkyl substances contamination in the livers of broiler chickens raised in Italy. *Ital J Food Saf* 14:13520.
- Domingo JL, Nadal M, 2017. Per- and polyfluoroalkyl substances (PFASs) in food and human dietary intake: a review of the recent scientific literature. *J Agric Food Chem* 65:533-43.
- EFSA, 2012a. Guidance on the Use of Probabilistic Methodology for Modelling Dietary Exposure to Pesticide Residues. *EFSA J* 10:2839.
- EFSA, 2012b. Guidance on selected default values to be used by the EFSA Scientific Committee, Scientific Panels and Units in the absence of actual measured data. *EFSA J* 10:2579.
- EFSA, 2020a. Risk to human health related to the presence of perfluoroalkyl substances in food. *EFSA J* 18:e06223.
- EFSA, 2020b. Human risk assessment of multiple chemicals using component-based approaches: a horizontal perspective. *EFSA J* 17:1759E.
- EFSA, 2025. Guidance on the use of read-across for chemical safety assessment in food and feed. *EFSA J* 23:e9586.
- European Chemicals Agency, 2016. How to use alternatives to animal testing to fulfil the information requirements for REACH registration. Practical guide. Available from: <https://data.europa.eu/doi/10.2823/194297>.
- European Commission, 2022. Directive proposal of the European Parliament and of the council amending Directive 2000/60/EC establishing a framework for Community action in the field of water policy, Directive 2006/118/EC on the protection of groundwater against pollution and deterioration and Directive 2008/105/EC on environmental quality standards in the field of water policy. Available from: <https://eur-lex.europa.eu/legal-content/EN/TXT/?uri=celex:52022PC0540>.
- European Parliament, Council of the European Union, 2020. Directive (EU) 2020/2184 of the European Parliament and of the Council of 16 December 2020 on the quality of water intended for human consumption. In: *Official Journal*, L 435/1, 23/12/2020.
- Gazzotti T, Sirri F, Ghelli E, Zironi E, Zampiga M, Pagliuca G, 2021. Perfluoroalkyl contaminants in eggs from backyard chickens reared in Italy. *Food Chem* 362:130178.
- Genualdi S, Beekman J, Carlos K, Fisher CM, Young W, DeJager L, Begley T, 2022. Analysis of per- and poly-fluoroalkyl substances (PFAS) in processed foods from FDA's total diet study. *Anal Bioanal Chem* 414:1189-99.
- Hossini H, Massahi T, Parnoon K, Nouri M, 2025. Per- and polyfluoroalkyl substances (PFAS) in milk and dairy products: a literature review of the occurrence, contamination sources, and health risks. *Food Addit Contam Part A* 42:1284-96.
- Lasters R, Groffen T, Eens M, Coertjens D, Gebbink WA, Hofman J, Bervoets L, 2022. Home-produced eggs: An important human exposure pathway of perfluoroalkylated substances (PFAS). *Chemosphere* 308:136283.
- Nobile M, Arioli F, Curci D, Ancillotti C, Scanavini G, Chiesa LM, et al, 2023. Incidence of perfluoroalkyl substances in commercial eggs and their impact on consumer's safety. *Foods* 12:3846.
- Nobile M, Curci D, Rampazzo G, Chiesa LM, Gazzotti T, Ghidini S, Arioli F, Panseri S, 2024. Per- and polyfluoroalkyl substances in blue crabs from the Adriatic Sea and consumer safety evaluation. *Int J Food Sci Technol* 59:7229-39.
- OECD, 2021. Reconciling terminology of the universe of per- and polyfluoroalkyl substances: recommendations and practical guidance. Available from: https://www.oecd.org/en/publications/reconciling-terminology-of-the-universe-of-per-and-polyfluoroalkyl-substances_e458e796-en.html.

- Piva E, Fais P, Ioime P, Forcato M, Viel G, Cecchetto G, Pascali JP, 2023. Per- and polyfluoroalkyl substances (PFAS) presence in food: comparison among fresh, frozen and ready-to-eat vegetables. *Food Chem* 410:135415.
- Rampazzo G, Arioli F, Pagliuca G, Depau G, Zironi E, Gazzotti T, 2025. Cumulative exposure and health risk assessment of PFAS in animal-derived foods using the relative potency factor approach. *Toxics* 13:931.
- Zahm S, Bonde JP, Chiu WA, Hoppin J, Kanno J, Abdallah M, Blystone CR, Calkins MM, Guang-Dong H, Dorman DC, Fry R, Guo H, Haug LS, Hofmann JN, Iwasaki M, Machala M, Mancini FR, Maria-Engler SS, Møller P, Ng JC, Pallardy M, Post GB, Salihovic S, Schlezinger J, Soshilov A, Steenland K, Steffensen IL, Tryndyak V, White A, Woskie S, Fletcher T, Ahmadi A, Ahmadi N, Benbrahim-Tallaa L, Bijoux W, Chittiboyina S, de Conti A, Facchin C, Madia F, Mattock H, Merdas M, Pasqual E, Suonio E, Viegas S, Zupunski L, Wedekind R, Schubauer-Berigan MK, 2024. Carcinogenicity of perfluorooctanoic acid and perfluorooctanesulfonic acid. *Lancet Oncol* 25:16-7.
- Zeilmaker MJ, Fragki S, Verbruggen EMJ, Bokkers BGH, Lijzen JPA, 2018. Mixture exposure to PFAS: a relative potency factor approach. Available online: <https://www.rivm.nl/bibliotheek/rapporten/2018-0070.pdf>

Table 1. 95th percentile lower bound of per- and polyfluoroalkyl substances concentration data in eggs and egg products extracted from Annex A of the EFSA 2020 opinion (EFSA 2020a).

Compound	P95 LB (ng/kg)
PFOA	36.0
PFOS	120.0

P95, 95th percentile; LB, lower-bound.

Table 2. European egg and egg products mean consumption and standard body weight used for each consumer group.

Consumer group	BW (kg)	Consumption (g/day)
Toddlers	12	10.1
Adolescents	45	15.9
Adults	70	18.6
Elderly	70	18.3

BW; body weight.

Table 3. Relative potency factors applied based on liver toxicity as a toxicological endpoint. Source: Bil *et al.*, (2021).

Compound	RPF	
	min	max
PFBA	0.1	
PFBS	0	
PFDA	4	10
PFDS	2	
PFDoDA	3	
PFHpA	0	1
PFHpS*	0.6	2
PFPeS*	0.001	0.6
PFHxA	0	
PFHxS	0.6	
PFHxDA	0.02	
PFNA	10	
PFOA	1	
PFOS	2	
PFODA	0.02	
PFPeA	0	
PFTeDA	0.3	
PFTTrDA*	0.3	3
PFUnDA	4	

*Based on read across (Bil *et al.*, 2021)

RPF, relative potency factor.

Table 4. PFOA equivalent intakes through eggs and egg products.

Consumer group	PEQ* (ng/kg bw per week)		
	PFOA	PFOS	Cumulative EWI**
Toddlers	0.21	1.42	1.63
Adolescents	0.09	0.60	0.69
Adults	0.07	0.46	0.53
Elderly	0.06	0.43	0.50

PEQ, PFOA equivalents; EWI, estimated weekly intake.