

Myokines in Home-Based Functional Electrical Stimulation-Induced Recovery of Skeletal Muscle in Elderly and Permanent Denervation

Sascha Sajer (1), Giulio Sauro Guardiero (2,3), Bianca Maria Scicchitano (4)

(1) Department of Physiko&Rheuma-Therapie, Institute for Physical Medicine, St. Pölten, Austria; (2) A&C M-C Foundation for Translational Myology, Padova, Italy; (3) Interdepartmental Research Centre of Myology, University of Padova, Italy; (4) Istituto di Istologia ed Embriologia, Università Cattolica del Sacro Cuore, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

This article is distributed under the terms of the Creative Commons Attribution Noncommercial License (CC BY-NC 4.0) which permits any noncommercial use, distribution, and reproduction in any medium, provided the original author(s) and source are credited.

Abstract

Neuromuscular disorders, disuse, inadequate nutrition, metabolic diseases, cancer and aging produce muscle atrophy and this implies that there are different types of molecular triggers and signaling pathways for muscle wasting. Exercise and muscle contractions may counteract muscle atrophy by releasing a group of peptides, termed myokines, to protect the functionality and to enhance the exercise capacity of skeletal muscle. In this review, we are looking at the role of myokines in the recovery of permanent denervated and elderly skeletal muscle tissue. Since sub-clinical denervation events contribute to both atrophy and the decreased contractile speed of aged muscle, we saw a parallel to spinal cord injury and decided to look at both groups together. The muscle from lifelong active seniors has more muscle bulk and more slow fiber-type groupings than those of sedentary seniors, demonstrating that physical activity maintains slow motoneurons that reinnervate the transiently denervated muscle fibers. Furthermore, we summarized the evidence that muscle degeneration occur with irreversible *Conus* and *Cauda Equina* syndrome, a spinal cord injury in which the human leg muscles may be permanently disconnected from the peripheral nervous system. In these patients, suffering with an extreme case of muscle disuse, a complete loss of muscle fibers occurs within five to ten years after injury. Their recovered tetanic contractility, induced by home-based Functional Electrical Stimulation, can restore the muscle size and function in compliant Spinal Cord Injury patients, allowing them to perform electrical stimulation-supported stand-up training. Myokines are produced and released by muscle fibers under contraction and exert both local and systemic effects. Changes in patterns of myokine secretion, particularly of IGF-1 isoforms, occur in long-term Spinal Cord Injury persons and also in very aged people. Their modulation in Spinal Cord Injury and late aging are also key factors of home-based Functional Electrical Stimulation - mediated muscle recovery. Thus, Functional Electrical Stimulation should be prescribed in critical care units and nursing facilities, if persons are unable or reluctant to exercise. This will result in less frequent hospitalizations and a reduced burden on patients' families and public health services.

Key Words: Human denervated muscle; SCI; Elderly, Recovery of external-work contractility; Home Based Functional Electrical Stimulation, Myokines, IGF-1.

Eur J Transl Myol 28 (4): 337-345, 2018

During aging, several morpho-functional changes occur in skeletal muscle, i.e., the generalized loss of muscle mass, reduced myofiber size, and the progressive reduction in muscle strength, leading to a pathologic condition known as sarcopenia and dynapenia.¹⁻³ This progressive age-related muscle wasting process is associated with an increased prevalence of falls, a

greater incidence of diseases and the loss of functional independence.^{4,5} Inadequate nutrition, disuse, neuromuscular disorders, and metabolic diseases are among the many possible causes for the age-related decline in skeletal muscle mass. Several interventions have been suggested to counteract sarcopenia, that imply different types of molecular triggers and signaling

pathways for muscle wasting.^{6,7} In particular, exercise and muscle contractions, whether volitional or induced by Electrical Stimulation (ES), may counteract muscle atrophy by releasing a group of peptides, termed myokines, to protect the functionality and to enhance the exercise capacity of skeletal muscle.⁸⁻¹⁰ This review summarizes the potential positive effects of exercise-induced myokines looking at the role of myokines and of other molecular mechanisms in the recovery of denervated and elderly muscle tissue.

Muscle plasticity after Aging or Spinal Cord Injury and home based Functional Electrical Stimulation

In recent years, we have performed basic research and clinical trials on the use of electrostimulation to provoke positive muscle plasticity adaptations after Spinal Cord Injury (SCI). These studies investigated the use of long impulse electrical stimulation as a treatment for long-term permanently denervated human muscles, i.e., home-based Functional Electrical Stimulation (hbFES).^{11,12} Purpose developed electrical stimulators and very large electrodes that are now both commercially available (<https://www.schuhfried.com/umbraco/Surface/AuthenticationSurface/Login?returnUrl=%2Fportal>) were designed and implemented to achieve our goals.^{13,14} Those studies demonstrated a tremendous improvement in muscle trophism and an increase in external muscle power.¹³⁻¹⁹ Muscle strength was indeed improved to the point that individuals could stand up with assistance from direct electrical stimulation of the denervated muscles as well as perform in-place stepping and walk 100-200 m. The hbFES rehabilitation program was initiated relatively late after denervation (more than 8 month after SCI) because of clinical constraints and/or the belief that electrical stimulation of denervated muscles is ineffective and may actually interfere with eventual myofiber reinnervation. These ideas are still maintained by some experts despite recent evidence that electrostimulation may indeed enhance nerve growth and appropriate muscle reinnervation.²⁰⁻²² Alternative options will hopefully be realized with the future stem cells approaches.²³ Management based on *in vivo* protocols, such as the induction of muscle regeneration by injection of anesthetics in local anoxic conditions or ex-vivo techniques such as the proliferation of autologous myoblasts derived from patient's muscle biopsies, is still to be fully explored by pre-clinical and clinical research.²⁴⁻²⁸ Although myokines will certainly have a main role, we leave to the experts in the field discussion of these approaches. The aim of the present review is to contribute the experience of muscle specialists (biologists, physiologists, and rehabilitation experts) to first the impact of denervation-reinnervation processes on shaping aging muscle fibers and then to the role of myokines in the overall process of muscle aging and in its effective countermeasures. Ten years of

experience in hbFES for SCI patients^{13,14} were translated with positive results in trials for rejuvenation of skeletal muscles of elderly persons.²⁹⁻³⁸ Myokines are produced and released by muscle fibers during contraction exercise and exert both local and systemic effects.¹⁰ Changes in patterns of myokines secretions during hbFES-induced muscle recovery in both long-term SCI persons and old people were observed, particularly with Insulin-like Growth factor 1 (IGF-1), suggesting that they are key factors of muscle recovery mediated by hbFES.

Role of myokines in volitional physical activity and hbFES to counteract muscle aging decay

The scientific literature on the aging nervous system is immense, in particular, the works on the decay of cognition and mobility. However, the impact of muscle denervation on aging skeletal muscle fibers is a relatively orphan topic.³⁹⁻⁴² This is mainly related to the undetermined molecular nature of the trophic factors released by motor neurons to the muscle fibers of the different types of motor units.⁴³ It is well known that such mechanisms contribute to neuromuscular junction development and maintenance; however, if and what chemical trophic factors influence the synchronized expression of the hundreds of nuclei belonging to a single muscle fiber remains a subject of hypotheses, while the synchronized spread of muscle action potential seems to be a more rational mechanism.⁴³ Two reviews provide an excellent summary of what is known or hypothesized about this subject.^{1,44} It is well established that physical exercise and dietary proteins, as well as the intake of specific amino acids, are able to counteract the processes related to the progression of muscle mass loss. This may have beneficial effects on improving the anabolic response of muscle in the elderly.⁴⁵ However, the world record series of Master athletes shows an almost linear decay in muscle power that starts at age 30 years and declines to zero, theoretically at approximate 110 years of age. It is worth stressing that this occurs despite the fact that these record-holding sportsmen are the best that the human species can provide at any time point in terms of both genetic sport talents and the capacity and opportunity to train with the best possible advice.⁴⁶ A series of investigations on counteracting muscle decay involving active or sedentary seniors and SCI patients can be found in which myokines play the main role in explaining the results after Electrical Stimulation. The studies show that volitional exercise and h-bFES are crucial for maintaining both motor neurons and muscle fibers,^{20-22,47} despite some doubts present in the literature.⁴⁸⁻⁵⁰ In addition to a progressive loss of muscle mass, aging presents also a conspicuous reduction in myofiber plasticity with alterations in muscle-specific transcriptional mechanisms.^{1,30,31} During the aging process, protein synthetic rates decrease and an increase in protein degradation follows, affecting characteristics

of muscle fibers.⁵¹ Recently, it was demonstrated that age-associated loss of OPA1 in muscle impacts muscle mass, metabolic homeostasis, systemic inflammation, and epithelial senescence.⁵² It is generally accepted that the failure to repair damage is a contributory cause of functional impairment with aging,^{8,9,53,54} and also promotes the detrimental replacement of functional contractile muscle with fibrous tissue.⁵¹ Volitional physical exercise can reverse these damaging processes in both aging and cancer cachexia.^{53,57-60} Interestingly, it has been shown that both acute and prolonged resistance exercise stimulates the proliferation of satellite cells in healthy sedentary subjects, though blunted in elderly people.^{53,57-60} The increased levels of myostatin,⁶¹ a negative regulator of muscle mass, may explain this fact.^{62,63} An increase in autophagy in the muscle of athletic people has been reported,⁶⁴⁻⁶⁷ suggesting that exercise may activate an important system that detoxifies muscle cells. Another major factor that is associated with physical exercise is IGF-1.⁶⁸ Its production by muscle increases after 5–10 min of moderate to high-intensity exercise.^{69,70} The evidence suggests that training and regular exercise modulate expression of myokines (e.g., IL-6) and IGF-1, thereby regulating functional autophagy, and thus increasing muscle strength and attenuating sarcopenia.^{71,72} There is strong evidence that IGF-1 secretion by muscle fibers also influences synaptogenesis in the brain,⁷³ and at the neuromuscular junctions.^{74,75} Further, agrin (along with other proteins such as MuSK, rapsyn, ChAT, Hb9, munc18, etc.) is important to denervation/reinnervation processes.^{76,77} Because elderly people are often reluctant or unable to participate adequately in physical exercise,⁷⁸ an alternative approach with Electrical Stimulation was designed and implemented. Krenn et al.,⁷⁹ designed a stimulator that especially addresses the requirements of elderly people. Subjects in the study were exposed to regular neuromuscular ES training for a period of 9 weeks. The outcome was an increase in muscle strength, associated with an increase in abundance of fast fibers, which indeed are the first to respond to ES and are related to the power of skeletal muscle.³⁰ In addition, with ES an increase in expression of IGF-1 factors as well as markers of both satellite cell proliferation and extracellular matrix remodeling was detected along with downregulation in the expression of proteases just as occurs during volitional physical exercise.³¹ Furthermore, Mosole et al.,⁸⁰ recently demonstrated that ES modulate also the Calcium (Ca^{2+})-handling proteins, NFAT, and related proteases. The aims were to study the molecular mechanisms that support functional muscle improvement by ES. Indeed, Ca^{2+} cycling and activation of specific molecular pathways are essential in contraction-induced muscle adaptation. This study attains human muscle sections and total homogenates prepared from biopsies obtained before (control) and after 9 weeks of training by ES on a group of elderly volunteers. Evidence for activation of

kinase/ phosphatase pathways after ES was obtained. Moreover, expression of Sarcalumenin, Calsequestrin, and sarco/endoplasmic reticulum Ca^{2+} -ATPase (Serca) isoforms was regulated by training. The conclusions were that neuromuscular ES applied to *Vastus lateralis* muscle of sedentary seniors combines fiber remodeling with activation of Ca^{2+} -Calmodulin molecular pathways and modulation of the key Ca^{2+} -handling proteins.⁸⁰ Collagen expression was also reported to be remodelled during both volitional physical exercise and ES; indeed, expression of three different forms of collagen was upregulated in electrically stimulated muscle.³⁰ However, the increase in collagen expression seems not to stimulate fibrosis as is shown by both morphological evidence and at the level of important fibrosis modulators, namely the increase in expression of miR29.⁷⁷ Several longitudinal studies have shown that regular exercise is beneficial to the aged population.^{1,82} The Interreg IVa project recruited therapy-stable sedentary seniors with a normal lifestyle who were trained for 9 weeks with either volitional exercise by leg press,⁸³ or Electrical Stimulation.^{29-31,77} Functional tests of trained subjects showed that LP and ES induced improvements in both leg muscle force and mobility tests.^{84,85} ES significantly increased the size of fast type muscle fibers, together with a significant increase of Pax7 and NCAM positive satellite cells. Furthermore, muscle biopsies did not present evidence of muscle damage and/or inflammation.³¹ Altogether, the results demonstrated that physical exercise, either voluntary or induced by ES, improves functional performance and the structural properties of aged muscles

Conclusive Remarks

Functional electrical stimulation using long biphasic impulses and large surface electrodes is able to restore muscle mass, force production and movement in humans even after years of complete irreversible denervation. Patients suffering with flaccid paraplegia (complete and permanent denervation of lower extremity muscles, e.g., *conus* and *cauda equina* syndrome) are especially good candidates for these approaches if hbFES managements start from 1 to 6 years after SCI.^{13,14} Electrical stimulation-induced up-regulation of myokines is an essential mechanism to achieve this goal. Furthermore, we recently demonstrated that two-years of hbFES also improve the thickness of the stimulated skin in *conus* and *cauda equina* syndrome patients, providing a valuable anti-decubitus effect.^{86,87} Thus, pilot human studies and the application of existing experimental knowledge support the use of h-bFES to improve the condition of long-term denervated muscles.⁸⁸ Taking the idea 2 steps further, in the future, we may thereby prepare the muscle in SCI patients for bridging and reconnecting it to the central motor control unit. Neither volitional training nor hbFES may stop the aging process as we have learned from the Master Athletes.⁴⁶ However, detrained elderly

persons may significantly improve their muscle structure and function to a level that allows safer and increased mobility. Our results show that ES is a safe home-based method able to counteract atrophy of fast type muscle fibers, a biomarker of skeletal muscle aging in sedentary seniors.^{30,31} If seniors are unable or reluctant to perform a volitional physical activity, the electrical stimulation-induced up-regulation of myokines is an important contributing mechanism. Adding h-bFES to good nutrition and exercise where possible (e.g., volitional Full Body In-Bed Gym)^{12,89,90} will allow the elderly and SCI patients (even those that will be stimulated by the recently developed soft subdural implants for the delivery of electrochemical neuromodulation therapies to the spinal cord)⁹¹⁻¹⁰³ to look forward to increased autonomy and quality of life, and the prospect of better professional and social integration. In case of severe muscle atrophy in the older elderly, commercially available neuromuscular electrical stimulators may improve mobility with substantial reductions of fall risk and of the severity of secondary medical problems. Thus, hbFES should be prescribed in critical care units and nursing facilities, if persons are unable or reluctant to exercise. This will result in less frequent hospitalizations and a reduced burden on patients' families and public health services.

List of acronyms

ES - Electrical Stimulation
hbFES – acute respiratory distress syndrome
IGF-1 - Insulin-like Growth factor 1
SCI - Spinal Cord Injury

Author's contributions

SS, GSG, and BMS conceived the review, analyzed the data and wrote the paper.

Acknowledgments

We thank all the persons that take part to the research activities, in particular the volunteers, which trained by hbFES

Funding: Supported by A&CM-C Foundation for Translational Myology; The Italian Ministero per l'Università e la Ricerca Scientifica e Tecnologica (M.U.R.S.T.); The EU Commission Shared Cost Project RISE (Contract no. QLG5-CT-2001-02191), Austrian Research Promotion Agency, Federal Ministry for Transport, Innovation and Technology (bmvit) and the Ludwig Boltzmann Society, Vienna, Austria; European Regional Development Fund - Cross Border Cooperation Program Slovakia -.Austria 2007–2013 (Interreg-IVa), project Mobilität im Alter, MOBIL, N_00033 (partners: Ludwig Boltzmann Institute of Electrical Stimulation and Physical Rehabilitation, Austria, and Faculty of Physical Education and Sports, Comenius University in Bratislava, Slovakia); Austrian national cofinancing of the Austrian Federal Ministry of Science and Research. Progetto di ricerca d'interesse di

Ateneo, Università Cattolica del Sacro Cuore, Linea D.3.2 2015 to BMS. The contents of this article are solely the responsibility of the authors.

Conflict of Interest

The authors declare that the research was conducted in the absence of any commercial or financial relationships that could be construed as a potential conflict of interest.

Ethical Publication Statement

We confirm that we have read the Journal's position on issues involved in ethical publication and affirm that this report is consistent with those guidelines.

Corresponding Author

Bianca M. Scicchitano, Istituto di Istologia ed Embriologia, Università Cattolica del Sacro Cuore, Fondazione Policlinico Universitario A. Gemelli IRCCS, Rome, Italy

E-mail: biancamaria.scicchitano@unicatt.it

E-mail of co-authors

Sajer Sascha: office@drsajer.at

Sauro Guardiero Giulio: sauroguardiero@gmail.com

References

1. Mitchell WK, Williams J, Atherton P, et al. Sarcopenia, dynapenia, and the impact of advancing age on human skeletal muscle size and strength; a quantitative review. *Front Physiol* 2012 Jul 11;3:260. doi: 10.3389/fphys.2012.00260. eCollection 2012.
2. Buonocore D, Rucci S, Vandoni M, et al. Oxidative system in aged skeletal muscle. *Muscles Ligaments Tendons J* 2012;1:85-90. Print 2011 Jul.
3. Cruz-Jentoft AJ, Baeyens J, Bauer JM, et al. Sarcopenia: European consensus on definition and diagnosis: Report of the European Working Group on Sarcopenia in Older People. *Age Ageing* 2010;39: 412-23.
4. Ryall JG, Schertzer JD, Lynch GS. Cellular and molecular mechanisms underlying age-related skeletal muscle wasting and weakness. *Biogerontology* 2008;9:213-28.
5. Rolland Y, Czerwinski S, Abellan Van Kan G, et al. Sarcopenia: its assessment, etiology, pathogenesis, consequences and future perspectives. *J Nutr Health Aging* 2008;12:433-50.
6. Sandri M, Sandri C, Gilbert A, et al. Foxo transcription factors induce the atrophy-related ubiquitin ligase atrogin-1 and cause skeletal muscle atrophy. *Cell* 2004;117:399-412.
7. Jackman RW, Kandarian SC. The molecular basis of skeletal muscle atrophy. *Am J Physiol. Cell*

Role of Myokines in muscle recovery

Eur J Transl Myol 28 (4): 337-345, 2018

- Physiol 2004;287:C834-43. 10.1152/ajpcell.00579.2003
8. Carosio S, Berardinelli MG, Aucello M, Musarò A. Impact of ageing on muscle cell regeneration. *Ageing Res Rev* 2009;10:35-42. doi: 10.1016/j.arr.2009.08.001. Epub 2009 Aug 13.
 9. Vinciguerra M, Musarò A, Rosenthal N. Regulation of muscle atrophy in aging and disease. *Adv Exp Med Biol* 2010;694:211-33.
 10. Leal LG, Lopes MA, Batista ML Jr. Physical Exercise-Induced Myokines and Muscle-Adipose Tissue Crosstalk: A Review of Current Knowledge and the Implications for Health and Metabolic Diseases. *Front Physiol* 2018;9:1307. doi: 10.3389/fphys.2018.01307.
 11. Kern H, Carraro U. Home-Based Functional Electrical Stimulation for Long-Term Denervated Human Muscle: History, Basics, Results and Perspectives of the Vienna Rehabilitation Strategy. *Eur J Transl Myol* 2014 Mar 27;24(1):3296. doi: 10.4081/ejtm.2014.3296. eCollection 2014 Mar 31.
 12. Carraro U, Gava K, Musumeci A, et al. Safe Antiaging Full-Body In-Bed Gym and FES for Lazy Persons: Home In-Bed Exercises for Fighting Muscle Weakness in Advanced Age, in *Rehabilitation Medicine for Elderly Patients*, eds S. Masiero and U. Carraro (Springer-Nature Book), 2018, pag. 43-52. ISBN 978-3-319-57405-9 ISBN 978-3-319-57406-6 (eBook) 30.
 13. Kern H, Carraro U, Adami N, et al. Home-based functional electrical stimulation rescues permanently denervated muscles in paraplegic patients with complete lower motor neuron lesion. *Neurorehabil Neural Repair* 2010;24:709-21. doi: 10.1177/1545968310366129. Epub 2010 May 11.
 14. Kern H, Carraro U, Adami N, et al. One year of home-based daily FES in complete lower motor neuron paraplegia: recovery of tetanic contractility drives the structural improvements of denervated muscle. *Neurol Res* 2010;32:5-12. doi: 10.1179/174313209X385644.
 15. Kern H, Boncompagni S, Rossini K, et al. Long-term denervation in humans causes degeneration of both contractile and excitation-contraction coupling apparatus that can be reversed by functional electrical stimulation (FES). A role for myofiber regeneration? *J Neuropathol Exp Neurol* 2004;63:919-31.
 16. Kern H, Rossini K, Carraro U, et al. Muscle biopsies show that FES of denervated muscles reverses human muscle degeneration from permanent spinal motoneuron lesion. *J Rehabil Res Dev* 2005;42:43-53.
 17. Carraro U, Rossini K, Mayr W, Kern H. Muscle fiber regeneration in human permanent lower motoneuron denervation: relevance to safety and effectiveness of FES-training, which induces muscle recovery in SCI subjects. *Artif Organs* 2005;29:187-91.
 18. Kern H, Salmons S, Mayr W, et al. Recovery of long-term denervated human muscles induced by electrical stimulation. *Muscle Nerve* 2005;31:98-101.
 19. Boncompagni S, Kern H, Rossini K, et al. Structural differentiation of skeletal muscle fibers in the absence of innervation in humans. *Proc Natl Acad Sci U S A* 2007;104:19339-44.
 20. Zealea D L, Rodriguez RJ, Kenny T, et al. Electrical stimulation of a denervated muscle promotes selective reinnervation by native over foreign motoneurons. *J Neurophysiol* 2002;87:2195-9.
 21. Willand MP. Electrical Stimulation Enhances Reinnervation After Nerve Injury. *Eur J Transl Myol* 2015;25, 243-48.
 22. Willand MP, Rosa E, Michalski B, et al. Electrical muscle stimulation elevates intramuscular BDNF and GDNF mRNA following peripheral nerve injury and repair in rats. *Neuroscience* 2016;334:93-104. doi: 10.1016/j.neuroscience.2016.07.040. Epub 2016 Jul 29.
 23. Huang H, Sun T, Chen L, et al. Consensus of clinical neurorestorative progress in patients with complete chronic spinal cord injury. *Cell Transplant* 2014;23, Suppl 1:S5-17. doi: 10.3727/096368914X684952. Epub 2014 Oct 9.
 24. Mussini I, Favaro G, Carraro U. Maturation, dystrophic changes and the continuous production of fibers in skeletal muscle regenerating in the absence of nerve. *J Neurophatol Exp Neurol* 1987;46:315-31.
 25. Carraro U, Dalla Libera L, Catani C. Myosin light and heavy chains in muscle regenerating in absence of the nerve: transient appearance of the embryonic light chains. *Exp Neurol* 1983;79:106-17.

Role of Myokines in muscle recovery

Eur J Transl Myol 28 (4): 337-345, 2018

26. Carraro U, Rossini K, Zanin ME, et al. Induced myogenesis in long-term permanent denervation: perspective role in Functional Electrical Stimulation of denervated legs in humans. *Basic Appl Myol* 2002;12:53-63.
27. Rossini K, Zanin ME, Podhorska-Okolow M, Carraro, U. To stage and quantify regenerative myogenesis in FES-Induced Functional Recovery of Human Long-Term Permanent Denervated Muscle. *Basic Appl Myol* 2002;20:277-86.
28. Yablonka-Reuveni Z. The skeletal muscle satellite cell: still young and fascinating at 50. *J. Histochem Cytochem* 2011;59:1041-59. doi: 10.1369/0022155411426780.
29. Kern H, Barberi L, Löfler S, et al. Electrical stimulation counteracts muscle decline in seniors. *Front Aging Neurosci* 2014 Jul 24;6:189. doi: 10.3389/fnagi.2014.00189. eCollection 2014.
30. Zampieri S, Mosole S, Löfler S, et al. Physical Exercise in Aging: Nine Weeks of Leg Press or Electrical Stimulation Training in 70 Years Old Sedentary Elderly People. *Eur J Transl Myol* 2015;25(4):237-242. doi: 10.4081/ejtm.2015.5374. eCollection 2015 Aug 24. Review.
31. Zampieri S, Pietrangelo L, Loeffler S, et al. Lifelong Physical Exercise Delays Age-Associated Skeletal Muscle Decline. *J Gerontol A Biol Sci Med Sci* 2015;70: 63-73. doi: 10.1093/gerona/glu006. Epub 2014 Feb 18.
32. Zampieri S, Mammucari C, Romanello V, et al. Physical exercise in aging human skeletal muscle increases mitochondrial calcium uniporter expression levels and affects mitochondria dynamics. *Physiol Rep* 2016;4:13005. pii: e13005. doi: 10.14814/phy2.13005.
33. Kern H, Hofer C, Loeffler S, et al. Atrophy, ultrastructural disorders, severe atrophy and degeneration of denervated human muscle in SCI and Aging. Implications for their recovery by Functional Electrical Stimulation, updated 2017. *Neurol Res* 2017;39:660-666. doi: 10.1080/01616412.2017.1314906. Epub 2017 Apr 13. Review.
34. Carraro U, Kern H, Gava P, et al. Recovery from muscle weakness by exercise and FES: lessons from Masters, active or sedentary seniors and SCI patients. *Aging Clin Exp Res* 2016;29:579-90.
35. Carraro U, Edmunds KJ, Gargiulo P. 3D false color computed tomography for diagnosis and follow-up of permanent denervated human muscles submitted to home-based Functional Electrical Stimulation. *Eur J Transl Myol* 2015;25:129-40. doi: 10.4081/ejtm.2015.5133. eCollection 2015 Mar 11.
36. Edmunds KJ, Gíslason MK, Arnadóttir ID, et al. Quantitative Computed Tomography and Image Analysis for Advanced Muscle Assessment. *Eur J Transl Myol* 2016;26:6015. doi: 10.4081/ejtm.2016.6015. eCollection 2016 Jun 13.
37. Edmunds K, Gíslason M, Sigurdsson S, et al. Advanced quantitative methods in correlating sarcopenic muscle degeneration with lower extremity function biometrics and comorbidities. *PLoS One* 2018;13(3):e0193241. doi: 10.1371/journal.pone.0193241. eCollection 2018.
38. Gargiulo P, Gíslason MK, Edmunds KJ, et al. CT-Based Bone and Muscle Assessment in Normal and Pathological Conditions. *Encyclopedia of Biomedical Engineering*, 2019; Vol. 2:119-34. <http://doi.org.10.1016/B978-0-801238-3.99920-3>.
39. Ling SM, Conwit RA, Ferrucci L, Metter EJ. Age-associated changes in motor unit physiology: observations from the Baltimore Longitudinal Study of Aging. *Arch Phys Med Rehabil* 2009;90:1237-40.
40. Mosole S, Carraro U, Kern H, et al. Long-term high-level exercise promotes muscle reinnervation with age. *J Neuropathol Exp Neurol* 2014;73:284-94. doi: 10.1097/NEN.0000000000000032.
41. Mosole S, Carraro U, Kern H, et al. Use it or Lose It: Tonic Activity of Slow Motoneurons Promotes Their Survival and Preferentially Increases Slow Fiber-Type Groupings in Muscles of Old Lifelong Recreational Sportsmen. *Eur J Transl Myol* 2016 Nov 25;26(4):5972. doi: 10.4081/ejtm.2016.5972. eCollection 2016 Sep 15.
42. Power GA, Dalton BH, Gilmore KJ, et al. Maintaining Motor Units into Old Age: Running the Final Common Pathway. *Eur J Transl Myol* 2017;27(1):6597. doi: 10.4081/ejtm.2017.6597. eCollection 2017 Feb 24.
43. Lømo T. The response of denervated muscle to long-term stimulation (1985, revisited here in 2014). *Eur J Transl Myol* 2014 Mar 27;24(1):3294. doi: 10.4081/ejtm.2014.3294. eCollection 2014 Mar 31.
44. Hepple RT, Rice CL. Innervation and neuromuscular control in ageing skeletal muscle. *J Physiol* 2015 Oct 6. doi: 10.1113/JP270561. [Epub ahead of print].

Role of Myokines in muscle recovery

Eur J Transl Myol 28 (4): 337-345, 2018

45. Scicchitano BM, Sica G. The Beneficial Effects of Taurine to Counteract Sarcopenia. *Curr Protein Pept Sci* 2018;19:673-80. doi: 10.2174/1389203718666161122113609.
46. Gava P, Kern H, Carraro U. Age-associated power decline from running, jumping, and throwing male masters world records. *Exp Aging Res* 2015;41:115-35. doi: 10.1080/0361073X.2015.1001648.
47. Gordon T, English AW. Strategies to promote peripheral nerve regeneration: electrical stimulation and/or exercise. *Eur J Neurosci* 2016;43:336-50. doi: 10.1111/ejn.13005. Epub 2015 Aug 14.
48. Donovan-Hall MK, Burridge J, Dibb B, et al. The views of people with spinal cord injury about the use of functional electrical stimulation. *Artif Organs* 2011;35:204-11. doi: 10.1111/j.1525-1594.2011.01211.x.
49. Hughes AM, Burridge JH, Demain SH, et al. Translation of evidence-based assistive technologies into stroke rehabilitation: users' perceptions of the barriers and opportunities. *BMC Health Serv Res* 2014;14:124. doi: 10.1186/1472-6963-14-124.
50. Bersch I, Tesini S, Bersch U, Frotzler A. Functional electrical stimulation in spinal cord injury: clinical evidence versus daily practice. *Artif Organs* 2015;39:849-54. doi: 10.1111/aor.12618.
51. Carnio S, LoVerso F, Baraibar MA, et al. Autophagy impairment in muscle induces neuromuscular junction degeneration and precocious aging. *Cell Rep* 2014;8:1509-21. doi: 10.1016/j.celrep.2014.07.061. Epub 2014 Aug 28.
52. Tezze C, Romanello V, Desbats MA, et al. Age-Associated Loss of OPA1 in Muscle Impacts Muscle Mass, Metabolic Homeostasis, Systemic Inflammation, and Epithelial Senescence. *Cell Metab* 2017;25:1374-89.e6. doi: 10.1016/j.cmet.2017.04.021. Epub 2017 May 25.
53. Barberi L, Scicchitano BM, De Rossi M, et al. Age-dependent alteration in muscle regeneration: the critical role of tissue niche. *Biogerontology* 2013;14:273-92. doi: 10.1007/s10522-013-9429-4. Epub 2013 May 12.
54. Barberi L, Scicchitano BM, Musaro A. Molecular and cellular mechanisms of muscle aging and sarcopenia and effects of electrical stimulation in seniors. 2015 *Eur J Transl Myol* 2009;25:231-6. doi: 10.4081/ejtm.2015.5227. eCollection 2015 Aug 24. Review.
55. Pigna E, Berardi E, Aulino P, et al. Aerobic Exercise and Pharmacological Treatments Counteract Cachexia by Modulating Autophagy in Colon Cancer. *Sci Rep* 2016 May 31;6:26991. doi: 10.1038/srep26991.
56. Coletti D. Chemotherapy-induced muscle wasting: an update. *Eur J Transl Myol* 2018 Jun 4;28(2):7587. doi: 10.4081/ejtm.2018.7587. eCollection 2018 Apr 24.
57. Musarò A. The Basis of Muscle Regeneration. *Advances in Biology* 2014. 2014;1-16.
58. Snijders T, Verdijk LB, van Loon LJ. The impact of sarcopenia and exercise training on skeletal muscle satellite cells. *Ageing Res Rev* 2009;8:328-38.
59. Mikkelsen UR, Langberg H, Helmark IC, et al. Local NSAID infusion inhibits satellite cell proliferation in human skeletal muscle after eccentric exercise. *J Appl Physiol* (1985) 2009;107:1600-1161. doi: 10.1152/jappphysiol.00707.2009. Epub 2009 Aug 27.
60. Farup J, Rahbek SK, Riis S, et al. Influence of exercise contraction mode and protein supplementation on human skeletal muscle satellite cell content and muscle fiber growth. *J Appl Physiol* (1985) 2014;117:898-909. doi: 10.1152/jappphysiol.00261.2014. Epub 2014 Aug 7.
61. Farup J, Rahbek SK, Knudsen IS, et al. Whey protein supplementation accelerates satellite cell proliferation during recovery from eccentric exercise. *Amino Acids* 2014;46:2503-26.
62. McKay BR, Ogborn DI, Bellamy LM, et al. Myostatin is associated with age related human muscle stem cell dysfunction. *FASEB J* 2012;26:2509-21. doi: 10.1096/fj.11-198663. Epub 2012 Mar 7.
63. McPherron AC, Lawler AM, Lee SJ. Regulation of skeletal muscle mass in mice by a new TGF-beta superfamily member. *Nature* 1997;387:83-90.
64. Kadi, F., Schjerling, P., Andersen, L. L., Charifi, N., Madsen, J. L., Christensen, L. R., et al. (2004). The effects of heavy resistance training and detraining on satellite cells in human skeletal muscles. *J. Physiol.* 2004;558:1005-12. Epub 2004 Jun 24.

Role of Myokines in muscle recovery

Eur J Transl Myol 28 (4): 337-345, 2018

65. Mackey AL, Holm L, Reitelseder S, et al. Myogenic response of human skeletal muscle to 12 weeks of resistance training at light loading intensity. *Scand J Med Sci Sports* 2010;21:773-82.
66. Scicchitano BM, Rizzuto E, Musarò A. Counteracting muscle wasting in aging and neuromuscular diseases: the critical role of IGF-1. *Aging (Albany NY)* 2009;1:451-7.
67. Sandri M, Barberi L, Bijlsma AY, et al. Signalling pathways regulating muscle mass in ageing skeletal muscle: the role of the IGF1-Akt- mTOR-FoxO pathway. *Biogerontology* 2013;14:303-23.
68. Adamo ML, Farrar RP. Resistance training, and IGF involvement in the maintenance of muscle mass during the aging process. *Ageing Res Rev* 2006;5:310-31. Epub 2006 Sep 1.
69. Wallace, JD, Cuneo RC, Baxter R, et al. Responses of the growth hormone (GH) and insulin-like growth factor axis to exercise, GH administration, and GH withdrawal in trained adult males: A potential test for GH abuse in sport. *J Clin Endocrinol Metab* 1999;84:3591-3601.
70. Kostka T, Patricot MC, Mathian B, et al. Anabolic and catabolic hormonal responses to experimental two-set low-volume resistance exercise in sedentary and active elderly people. *Aging Clin Exp Res* 2003;15:123-30.
71. Berg U, Bang P. Exercise and circulating insulin-like growth factor I. *Horm Res* 2004;62:50-8.
72. Kern H, Pelosi L, Coletto L, et al. Atrophy/hypertrophy cell signaling in muscles of young athletes trained with vibrational-proprioceptive stimulation. *Neurol Res* 2011;33:998-1009.
73. Pelosi L, Berardinelli MG, De Pasquale L. Functional and Morphological Improvement of Dystrophic Muscle by Interleukin 6 Receptor Blockade. *EBioMedicine* 2015;2: 274-5.
74. Nieto-Estévez V, Defterali Ç, Vicario-Abejón C. IGF-I: A Key Growth Factor that Regulates Neurogenesis and Synaptogenesis from Embryonic to Adult Stages of the Brain. *Front Neurosci* 2016;10:52. doi: 10.3389/fnins.2016.00052. eCollection 2016.
75. Apel PJ, Ma J, Callahan M, et al. Effect of locally delivered IGF-1 on nerve regeneration during aging: an experimental study in rats. *Muscle Nerve* 2010;41:335-41. doi: 10.1002/mus.21485.
76. Wang J, Jokerst JV. Stem Cell Imaging: Tools to Improve Cell Delivery and Viability. *Stem Cells Int* 2016;2016:9240652. doi: 10.1155/2016/9240652. Epub 2016 Jan 6.
77. Tezuka T, Inoue A, Hoshi T, et al. The MuSK activator agrin has a separate role essential for postnatal maintenance of neuromuscular synapses. *Proc Natl Acad Sci U S A* 2014;111:16556-61. doi: 10.1073/pnas.1408409111. Epub 2014 Nov 3.
78. Bily W, Franz C, Trimmel L, et al. Effects of Leg-Press Training With Moderate Vibration on Muscle Strength, Pain, and Function After Total Knee Arthroplasty: A Randomized Controlled Trial. *Arch. Phys. Med. Rehabil.* 2016;97:857-65. doi: 10.1016/j.apmr.2015.12.015. Epub 2016 Jan 4.
79. Krenn M, Haller M, Bijak, M, et al. Safe neuromuscular electrical stimulator designed for the elderly. *Artif Organs* 2011;35:253-6.
80. Mosole S, Zampieri S, Furlan S, et al. Effects of Electrical Stimulation on Skeletal Muscle of Old Sedentary People. *Gerontol Geriatr Med* 2018;4:2333721418768998. doi: 10.1177/2333721418768998. eCollection 2018 Jan-Dec.
81. He Y, Huang C, Lin, X, Li J. MicroRNA-29 family, a crucial therapeutic target for fibrosis diseases. *Biochimie* 2013;95:1355-9.
82. Carraro U, Franceschi C. Apoptosis of skeletal and cardiac muscles and physical exercise. *Aging (Milano)*. 1997;9:19-34. Review.
83. Hamar D. Universal linear motor driven Leg Press Dynamometer and concept of Serial Stretch Loading. *Eur J Transl Myol* 2015;25:215-9.
84. Cvecka J, Tirpakova V, Sedliak M, et al. Physical activity in elderly. *Eur. J. Transl. Myol.* 2015; 25:249-52.
85. Sarabon N, Löfler S, Hosszu G, Hofer Ch. Mobility test protocols for the elderly: a methodological note. *Eur J Transl Myol* 2015;25:253-6.
86. Albertin G, Kern H, Hofer C, et al. Two years of Functional Electrical Stimulation by large surface electrodes for denervated muscles improve skin epidermis in SCI. *Eur J Transl Myol* 2018;28:7373. doi: 10.4081/ejtm.2018.7373. eCollection 2018 Jan 12.
87. Albertin G, Hofer C, Zampieri S, et al. In complete SCI patients, long-term functional electrical stimulation of permanent denervated muscles increases epidermis thickness. *Neurol Res* 2018;40:277-82. doi:

- 10.1080/01616412.2018.1436877. Epub 2018 Feb 15.
88. Kern H, Gargiulo P, Pond A, et al. To reverse atrophy of human muscles in complete SCI lower motor neuron denervation by home-based Functional Electrical Stimulation, in *Muscle Atrophy*, ed J. Xiao. *Advances in Experimental Medicine and Biology* 2019;1088, https://doi.org/10.1007/978-981-13-1435-3_252019. https://doi.org/10.1007/978-981-13-1435-3_27.
 89. Carraro U, Gava K, Baba A, et al. Fighting muscle weakness in advanced aging by take-home strategies: Safe anti-aging full-body in-bed gym and functional electrical stimulation (FES) for mobility compromised elderly people. *Biol Eng Me* 2018;1: 1-4.
 90. Carraro U, Gava K, Baba A, et al. To contrast and reverse skeletal muscle atrophy by Full-Body In-Bed Gym, a mandatory life-style for older olds and borderline mobility impaired persons, in *Muscle Atrophy*, ed J. Xiao. *Adv Exp Med Biol* 2019;1088, https://doi.org/10.1007/978-981-13-1435-3_25.
 91. Ekelem A, Goldfarb M. Supplemental stimulation improves swing phase kinematics during exoskeleton assisted gait of SCI subjects with severe muscle spasticity. *Front Neurosci* 2018 Jun 1;12:374. doi: 10.3389/fnins.2018.00374. eCollection 2018.
 92. Angeli CA, Boakye M, Morton RA, et al. Recovery of Over-Ground Walking after Chronic Motor Complete Spinal Cord Injury. *N Engl J Med* 2018;379(13):1244-50. doi: 10.1056/NEJMoa1803588. Epub 2018 Sep 24.
 93. Harkema SJ, Legg Ditterline B, Wang S, et al. Epidural Spinal Cord Stimulation Training and Sustained Recovery of Cardiovascular Function in Individuals With Chronic Cervical Spinal Cord Injury. *JAMA Neurol* 2018 Sep 17. doi: 10.1001/jamaneurol.2018.2617. [Epub ahead of print].
 94. Herrity AN, Williams CS, Angeli CA, et al. Lumbosacral spinal cord epidural stimulation improves voiding function after human spinal cord injury. *Sci Rep* 2018 Jun 6;8(1):8688. doi: 10.1038/s41598-018-26602-2.
 95. Aslan SC, Legg Ditterline BE, Park MC, et al. Epidural Spinal Cord Stimulation of Lumbosacral Networks Modulates Arterial Blood Pressure in Individuals With Spinal Cord Injury-Induced Cardiovascular Deficits. *Front Physiol* 2018;9:565. doi: 10.3389/fphys.2018.00565. eCollection 2018.
 96. Harkema SJ, Wang S, Angeli CA, et al. Normalization of Blood Pressure With Spinal Cord Epidural Stimulation After Severe Spinal Cord Injury. *Front Hum Neurosci* 2018;12:83. doi: 10.3389/fnhum.2018.00083. eCollection 2018.
 97. Terson de Paleville DGL, Harkema SJ, Angeli CA. Epidural stimulation with locomotor training improves body composition in individuals with cervical or upper thoracic motor complete spinal cord injury: A series of case studies. *J Spinal Cord Med* 2018 Mar 14;1-7. doi: 10.1080/10790268.2018.1449373. [Epub ahead of print]
 98. Hubscher CH, Herrity AN, Williams CS, et al. Improvements in bladder, bowel and sexual outcomes following task-specific locomotor training in human spinal cord injury. *PLoS One* 2018;13(1):e0190998. doi: 10.1371/journal.pone.0190998. eCollection 2018.
 99. Wagner FB, Mignardot JB, Le Goff-Mignardot CG, et al. Targeted neurotechnology restores walking in humans with spinal cord injury. *Nature* 2018 Nov;563(7729):65-71. doi: 10.1038/s41586-018-0649-2. Epub 2018 Oct 31.
 100. Formento E, Minassian K, Wagner F, et al. Electrical spinal cord stimulation must preserve proprioception to enable locomotion in humans with spinal cord injury. *Nat Neurosci* 2018 Oct 31. doi: 10.1038/s41593-018-0262-6. [Epub ahead of print]
 101. Capogrosso M, Wagner FB, Gandar J, et al. Configuration of electrical spinal cord stimulation through real-time processing of gait kinematics. *Nat Protoc* 2018 Sep;13:2031-61. doi: 10.1038/s41596-018-0030-9.
 102. Moraud EM, von Zitzewitz J, Miehlebradt J, et al. Closed-loop control of trunk posture improves locomotion through the regulation of leg proprioceptive feedback after spinal cord injury. *Sci Rep* 2018;8:76. doi: 10.1038/s41598-017-18293-y

Received for publication 31/8/2018

Revision received: 06/11/2018

Accepted for publication: 08/11/2018