The antioxidant effects of vitamin C on liver enzymes: aspartate aminotransferase, alanine aminotransferase, alkaline phosphatase and gamma-glutamyltransferase activities in rats under Paraquat insult

Benjamin Nnamdi Okolonkwo,
Edna Ogechi Nwachuku

Department of Medical Laboratory
Sciences, Rivers State University
of Science and Technology,
Nkpolu-Orowuroko, Nigeria

Abstract

Paraquat (PQ) is a bipyridylium herbicide; applied around trees in orchards and between crop rows to control broad-leaved and grassy weeds. Its oxidation results in the formation of superoxides which causes damage to cellular components. In this study, we determined the antioxidant effect vitamin C has on the liver enzymes [aspartate aminotransferase (SGOT), alanine aminotransferase (SGPT), alkaline phosphatase (ALP), and gamma-glutamyltransferase (GGT)] of rats under this toxic insult. Male rats in groups (A, B, C and D) were intraperitoneally injected with different sub-lethal increasing doses (0, 0.02, 0.04 and 0.06 g/kg body weight) of PQ respectively on monthly basis. Subsequently, the subgroups (A_2 , B_2 , C_2 and D_2) were given orally, 200 mg/L vitamin C, while the subgroups A_1 , B_1 , C_1 , and D_1 , received only water. Four animals per subgroup were decapitated on monthly basis and blood samples taken for enzyme assay. The parameters studied were - SGOT, SGPT, ALP and GGT - liver enzymes. The dose and time dependent PQ toxicity effect resulted in highly elevated Liver enzymes activities. The subgroups on vitamin C had significantly lower enzyme activities when compared to the same subgroups on only PQ insult. But the values were high when compared to the control subgroups (A_1 and A_2). These results were indication that vitamin C when given at moderate doses and maintained for a longer period could be a life saving adjunct to toxic insult.

Introduction

Paraquat (PQ), a bipyridylium herbicide that is effective as non-selective herbicide

when applied to leaves.¹ Because Paraquat has a redox potential of -446 mv, any reducing agent with sufficient energy can donate an electron to the bipyridylium divalent cation, Paraquat²⁺, to form a free radical, paraquat[•]. The oxidation of the bipyridylium radical to form the original paraquat²⁺ results in the transfer of the electron to oxygen and the formation of superoxides.² Subsequent Haber-Weiss and Fenton reactions yield toxic hydroxyl radicals. Thus the herbicide (paraquat) functions as a catalyst to transfer reducing equivalent to oxygen. These reactive oxygen species (ROS) so formed may escape the electron transport chain and cause damage to cellular components.³

In general, Antioxidant systems either prevent these ROS from being formed, or remove them before they can damage vital components of the cell.⁴ Antioxidants are classified into two broad divisions, depending on their solubility in water (hydrophilic) or in lipids (hydrophobic or lipophilic). Water – soluble antioxidants reacts with oxidants in the cell cytosol and the blood plasma, while lipid – soluble ones protects cell membranes from lipid peroxidation.⁵ Though various antioxidants behave synergistically, ascorbic acid (vitamin C) can degrade other antioxidants or reactivate them.⁵ Vitamin C (ascorbate) can directly scavenge oxygen free radicals with and without enzyme catalysts and can indirectly scavenge them by recycling others to their reduced form.⁶ By reacting with activated oxygen more readily than any other aqueous components, vitamin C protects critical macromolecules from oxidative damage.⁶

The relative importance of vitamin C as an antioxidant is the reason this research was centered on how to enhance the antioxidant system in the body to counter the destructive effects of Paraquat. The liver enzyme was so chosen because the liver is the energy store and source of the body.

Materials and Methods

Rats

A total of 100 male albino rats (*Rattus norvegicus*), weighing between 180-220 g [average body weight (bw) of 0.2 ± 0.02 kg], obtained from the animal house of the Department of Pharmacology and Toxicology, College of Health Sciences, University of Port Harcourt, Choba, Rivers State, Nigeria, were fed *ad libitum* with animal pelletized finisher feed with negligible vitamin c content, and allowed to acclimatize for two weeks in metabolic cages before the commencement of the study.

Correspondence: Benjamin Nnamdi Okolonkwo, Department of Medical Laboratory Sciences, Faculty of Sciences, Rivers State University of Science and Technology, Nkpolu-Orowuroko, Port-Harcourt, Rivers State, Nigeria.
Tel. +234.805.614.1653.
E-mail: benbruceph@yahoo.com

Key words: Paraquat, antioxidant, vitamin C, reactive oxygen species.

Contributions: BNO, as the lead author, was involved in the research proper, analyzing the samples and making sure that all went well with the study; EON, carried out the statistical analysis and assisted in proof reading of the materials before publication.

Conflict of interests: the authors declare no potential conflict of interests.

Received for publication: 14 February 2013.

Revision received: 6 May 2013.

Accepted for publication: 14 May 2013.

This work is licensed under a Creative Commons Attribution NonCommercial 3.0 License (CC BY-NC 3.0).

©Copyright B.N. Okolonkwo and E.O. Nwachuku, 2013

Licensee PAGEPress, Italy

Journal of Xenobiotics 2013; 3:e5

doi:10.4081/xeno.2013.e5

Paraquat

A percentage of 20 w/v Dizmazone® (Paraquat solution) from Dizengoff W.A. Ltd. (Lagos, Nigeria) sealed in an opaque plastic container, with a shelf-life of two years.

Vitamin C

Four bottles of Mason Natural® - pure Vitamin C (100 mg) caplets were employed. They are products of Mason Vitamins, Inc. (Miami Lakes, FL, USA).

Method

Two mL of sub-lethal doses of the toxicant (PQ) was intraperitoneally (i.p.) administered to the animals, under anaesthetics,⁷ in different dosed subgroups – A_1 , A_2 (0 g/kg); B_1 , B_2 (0.02 g/kg); C_1 , C_2 (0.04 g/kg) and D_1 , D_2 (0.06 g/kg) – on bi-weekly basis over a period of three months (in simulation of contamination from polluted feed, water or air), while the control animals (subgrouped A_1 , A_2) received 2 mL of normal saline (0.98 %) likewise in conformity with international standard in checking injection site reaction. The subgrouped animals were designated into: i) for non vitamin C treatment; and ii) for vitamin C treatment as indicated in Table 1.

This study was conducted under eight sub-

groups (A₁, A₂, B₁, B₂, C₁, C₂, D₁ and D₂). All the subgroups had 12 animals each with A₁ and A₂ being the control subgroups that received no Paraquat, A₂, B₂, C₂ and D₂ were placed on vitamin C – they were made to drink from water bottles with glass sipper tubes that contained vitamin c solution (200 mg/L), A₁, B₁, C₁ and D₁ were the subgroups that received ordinary drinking water with negligible vitamin C content. The water and vitamin c solution bottles were refilled at least trice daily irrespective of the volume of water or vitamin c remaining in the in-use bottles. It will be note worthy to state that at month 3, the food and water consumption by the rats were affected by PQ intoxication, mainly the subgroups dosed 0.06 g/kg bw without vitamin c treatment (D₁).

On monthly intervals, four animals per subgroup were selected, anaesthetized with gaseous isoflurane anesthetic machine, the induction chamber was prefilled with 4% isoflurane and oxygen (0.6 L/min). The rats were placed in the induction chamber and observed for signs of lateral recumbence, (steady breathing and no attempt to right itself when the induction chamber is slightly tilted), only then it becomes anaesthetized enough for transfer to the mask on the rodent breathing circuit. Open the lid of the induction chamber and quickly check for absence of the pedal reflex,⁸ if present, 10 mL of blood sample were collected using cardiac puncture procedures.⁷

The samples were processed and centrifuge, with the serum separated and labeled accordingly; stored frozen until needed for the estimation of enzyme activity using Kodac Autoanalyser machine. Using the study pattern below, the results were as shown in Tables 1 and 2 and in Figures 1-4.

Animal care

We do affirm that in carrying out this

research that *The Nigerian Institutional and National Guide for the care and use of laboratory animals* were followed.

Data computation

The Excel (2007) window's package and two-way analysis of variance (ANOVA) statistical methods were used for the result analysis. With levels of significance measured at $P \leq 0.05$ and 0.001, respectively.

Table 1. Treatment chart for the subgroups.

Group A: Control group given 0.0 g/kg PQ and divided into two subgroups	
A1: subsequently received water and feed	
A2: subsequently received vitamin C solution (1000 mg/L)	
Group B: Test group given 0.02 g/kg PQ and divided into two subgroups	
B1: subsequently received water and feed	
B2: subsequently received vitamin C solution (1000 mg/L) and feed	
Group C: Test group given 0.04 g/kg PQ and divided into two subgroups	
C1: subsequently received water and feed	
C2: subsequently received vitamin C solution (1000 mg/L) and feed	
Group D: Test group given 0.04g/kg PQ and divided into two subgroups	
D1: subsequently received water and feed	
D2: subsequently received vitamin C solution (1000 mg/L) and feed	

PQ, Paraquat.

Table 2. Values of the enzymes activities within the 3 months.

Months	Rat no.	Group	Sub-group	SGOT (IU/L)	SGPT (IU/L)	ALP (IU/L)	GGT (IU/L)
1	4	A	A1	2.2±0.4	2.52±0.8	11.25±3.0	13.63±3.8
	4		A2	4.6±1.1	6.50±0.9	14.72±1.9	12.17±2.6
	4	B	B1	15.4±1.2	10.95±2.0	53.44±11.2	32.00±5.6
	4		B2	12.7±1.9	12.53±2.1	33.94±8.5	22.33±3.3
	4	C	C1	46.2±5.6	134.88±23.4	82.00±17.5	42.67±9.9
	4		C2	41.5±6.7	83.87±15.9	20.83±3.2	26.33±5.5
	4	D	D1	99.5±10.1	155.67±36.9	318.17±39.0	65.00±13.7
	4		D2	44.5±8.9	133.23±28.3	272.00±46.1	35.50±4.7
2	4	A	A1	5.6±1.3	4.18±2.7	15.70±0.5	13.33±1.7
	4		A2	6.2±1.2	6.25±2.4	7.38±2.8	13.51±7.0
	4	B	B1	19.8±4.7	41.15±4.4	100.30±10.0	63.00±13.9
	4		B2	15.1±1.3	31.14±6.3	71.74±15.4	36.88±3.0
	4	C	C1	94.7±11.8	147.00±15.2	130.68±8.4	76.25±13.0
	4		C2	74.3±13.0	96.25±11.8	82.33±27.6	35.13±2.2
	4	D	D1	138.5±22.3	258.75±31.9	315.53±16.34	99.00±6.0
	4		D2	107.6±18.9	142.00±34.2	185.18±6.5	76.00±11.3
3	4	A	A1	4.2±0.9	1.35±0.6	19.60±0.2	12.75±0.8
	4		A2	6.8±3.4	7.50±1.2	13.53±1.5	11.60±1.8
	4	B	B1	54.1±8.0	99.00±1.0	1028.00±44.2	87.00±13.0
	4		B2	48.3±5.5	46.50±3.5	527.50±35.1	50.00±21.0
	4	C	C1	115.2±20.0	191.00±24.0	1297.00±47.9	128.50±8.5
	4		C2	101.5±5.5	74.50±10.5	691.00±55.8	56.50±3.5
	4	D	D1	342.0±18.7	167.50±12.5	1570.00±68.0	170.00±6.4
	4		D2	199.0±18.1	58.00±6.0	1096±37.7	125.00±12.0

SGOT, aspartate aminotransferase; SGPT, alanine aminotransferase; ALP, alkaline phosphatase; GGT gamma-glutamyltransferase.

Results

The SGOT activity of all the subgroups is shown in Figure 1. The enzyme activity values were found to have increased on the test subgroups (B_1 , B_2 , C_1 , C_2 , D_1 and D_2) when compared to the control subgroups (A_1 and A_2). And the increases were found to be directly related to the dose and exposure time effects ($P \leq 0.001$). As the time of exposure moved from month 1 down to month 3 the enzyme activity of the PQ only treated subgroups (B_1 , C_1 and D_1) increased two- to three- folds that of the subgroups that received vitamin C in addition to PQ insult (B_2 , C_2 and D_2) at the same level of significance ($P \leq 0.001$), indicating that an interaction exists between dose of PQ/vitamin C given and time of exposure. The test subgroups with the highest dose 0.06 g/kg bw (D_1 and D_2) gave an explicit picture on how vitamin C effected an improved enzyme activity when compared to the one on PQ insult only. With D_1 being higher than two-times D_2 at months 1 and 3.

The results of the SGPT enzyme activity values as shown in Figure 2 were similar to the ones obtained in SGOT (Figure 1) values above. There existed an interaction between dose given and time of exposure effects at $P \leq 0.001$ level of significance. The test subgroups (B_1 , B_2 , C_1 , C_2 , D_1 and D_2) were highly elevated when compared to the control subgroups (A_1 and A_2) from month 1 to month 3 especially the subgroups dosed 0.04 g/kg and 0.06 g/kg bw (C_1 , C_2 , D_1 and D_2 at month 1) and (B_1 , B_2 , C_1 , C_2 , D_1 and D_2 at months 2 and 3). The within group comparison indicated a well defined reduction in the values of the SGPT enzyme activity of the vitamin C treated subgroups (B_2 , C_2 and D_2) when compared to the subgroups on PQ insult only (B_1 , C_1 and D_1) from month 1 to month 3. This results was in line with the results of the SGOT enzyme activity values obtained in Figure 1, showing that vitamin C has an ameliorative effect on intracellular PQ toxicity.

The results obtained from ALP enzyme activity on the subgroups were quite remarkable. At month 1, the test subgroups with lower doses of PQ (B_1 , B_2 , C_1 and C_2) showed slight elevation of ALP activity when compared to the control subgroups (A_1 and A_2) at $P \leq 0.05$, while the subgroups that received the highest dose of 0.06g PQ/kg bw (D_1 and D_2) had an elevated ALP activity at $P \leq 0.01$ when compared to the control subgroups. As the time of exposure increased from month 1 to month 3 the level of enzyme activity of the test subgroups increased the more to a level that was quite high at month 3 ($P \leq 0.001$). In all this, we found out that the vitamin C treated test sub-

groups had lower ALP enzyme activity values than the subgroups under the same PQ insult only ($B_2 < B_1$, $C_2 < C_1$, $D_2 < D_1$) at $P \leq 0.05$ (month 1), $P \leq 0.01$ (month 2) and $P \leq 0.001$ (month 3) respectively. This result showed that an interaction existed between the dose of PQ given and time of exposure of the toxic insult, and also there was a relationship between the effect of the vitamin C given with the dose of PQ given and the time of exposure.

At month 1 the GGT enzyme activity values of the test subgroups were increasing in direct

relation to the increase in dose of the toxicant (PQ) at $P \leq 0.001$, and this relationship continued to increase the enzyme activity even as the exposure time increased from month 1 to month 3 at $P \leq 0.001$. Also the vitamin C treated subgroups (B_2 , C_2 and D_2) all had much lower enzyme activity values than their counterpart under PQ insult only (B_1 , C_1 and D_1) at $P \leq 0.001$ from month 1 through to month 3. This also supports the existence of an interaction between the dose of PQ given and time of exposure of the toxic insult, and also a rela-

Figure 1. Aspartate aminotransferase (SGOT enzyme) activity values for 3 months study period.

Figure 2. Alanine aminotranferase (SGPT enzyme) activity values for 3 months study period.

tionship between the effect of the vitamin c given with the dose of PQ given and the time of exposure.

Discussion

PQ studies and the toxicity effects of this chemical on the organs of the body, including its mechanism of toxicity have been reported.⁹ Furthermore, several reports presented PQ as a hepatotoxin.¹⁰⁻¹³

Following the findings from the acute toxicity study, where it was observed that toxicity effects existed with increasing dose and time of exposure of PQ (i.p.) insult on the liver cells.¹⁴ A

chronic study was designed to assess the toxicity of PQ at long duration (3 months) and also to find the possibility of ameliorating the toxicity effects using antioxidant (vitamin C).

From the liver enzymes – SGOT, SGPT, ALP and GGT – results obtained, all had changes that were highly significant ($P \leq 0.001$) both within and between subgroups (Table 2 and Figures 1-4), and these changes were all dose and exposure-time dependent. The enzymes activity values of the subgroups that received PQ only were almost two to five folds increased compared to that of the subgroups that received vitamin C in addition to the PQ insult. SGOT values of vitamin C treated subgroups were significantly lower than those on PQ insult only ($B_2 < B_1$, $C_2 < C_1$, and $D_2 < D_1$ from

month 1 to month 3 (Figure 1) except the control group A, in-which the vitamin C treated subgroup (A_2) had slightly elevated SGOT values throughout the three months than the subgroup without vitamin C and PQ (A_1) ($A_1 < A_2$). These changes were also observed in the values of other enzymes – SGPT, ALP and GGT (Figures 2-4).

It has been shown that PQ toxicity initiates lipid peroxidation which causes toxic destruction of lipid membrane bilayers initiating release of membrane bound enzymes – SGPT, ALP and GGT – to the cytoplasm.¹⁵⁻¹⁸ This explains the highly elevated values of these enzymes in PQ only treated subgroups (B_2 , C_1 and D_1) as compared to the subgroups that received, in addition to PQ insult, vitamin C (B_2 , C_2 , and D_2) in Figures 2-4. Indicating that vitamin C, to a large extent, reduced the toxic insults⁶ and acted as a substrate for the antioxidant enzyme ascorbate peroxidase, a function that is particularly important in stress resistance,¹⁹ thereby maintaining and repairing cellular integrity and function¹⁵⁻¹⁷ which led to the lowered values obtained in (B_2 , C_2 and D_2) as seen in Figures 2-4.

Generally, there was an improvement in enzyme activity in the subgroups on vitamin C during the PQ toxic insult (B_2 , C_2 , and D_2) in Figures 1-4. Though these values were still high when compared to the control subgroups (A_1 and A_2) and above the reference values for rat's liver enzymes, it's still a pointer to health improvement. These shows that if vitamin c treatment was continued for a longer period it could have totally repaired the liver cells and improve the health status of the animals.

Figure 3. Alkaline phosphatase (ALP enzyme) activity values for 3 months study period.

Figure 4. Gamma-glutamyltransferase (GGT enzyme) activity values for 3 months study period.

Conclusions

The result of this study has demonstrated that vitamin c has the capacity to improve the health status of animals under toxic insult. Exposure of rats to PQ induced a highly elevated liver enzyme – SGOT, SGPT, ALP and GGT – activities that were dose and time dependent, which on subsequent administration of vitamin C a reduction in the activity levels were observed throughout the study, indicating the reliability of vitamin c as an adjunct to the treatment of a toxic insult.

Recommendation

Vitamin C, a potent antioxidant should be one of the first line treatments (emergency procedure) for toxic insult. Vitamin C administration in PQ toxicity should not be ignored, and it should be extended even after patient's

recovery for adequate intracellular repairs to be achieved.

References

1. Mees GC. Experiments on the herbicidal action of 1, 1'-ethylene- 2, 2'- dipyridylum dibromide. *Ann Appl Biol* 1960;48:601-12.
2. Punchard NA, Kelly FJ. Free radicals: a practical approach. Oxford: IRL press; 1996.
3. Davies K. Oxidative stress: the paradox of aerobic life. *Biochem Soc Symp* 1995; 61:1-31.
4. Rhee SG. Cell signaling. H₂O₂, a necessary evil for cell signaling. *Science* 2006;312:1882-3.
5. Sies H. Oxidative stress: oxidants and antioxidants. *Exp. Physiol* 1997;82:291-5.
6. Asada K. Ascorbate peroxidase – hydrogen peroxide scavenging enzyme in plants. *Physiol Plant* 1992;85:235-41.
7. Animal Care and Ethics Committee. Blood collection in rodents and rabbits. Newcastle: University of Newcastle, ACEC; 1999. Available from: <http://www.newcastle.edu.au/research/animal/revision/acec29.pdf> Accessed: June 2007.
8. Diel KH, Hull R, Morton D, Pfister R, Rabemampianina Y, Smith D, et al. A good practice guide to the administration of substances and removal of blood including routes and volumes. *J Appl Toxicol* 2001; 21:15-23.
9. Bus JS, Cogen SZ, Olgaard M, Gibson JE. A mechanism of paraquat toxicity in mice and rats. *Toxicol Appl Pharmacol* 1976;35: 501-13.
10. Clark DG, McElligott TF, Hurst EW. The toxicity of paraquat. *Br J Ind Med* 1966;23: 126-32.
11. Gibson JE, Cagen SZ. Paraquat – induced functional changes in kidney and liver. In: Wong, RC, Stevens, JB, eds. *Biochemical mechanisms of paraquat toxicity*. 2nd ed. New York: Academic Press; 1977. pp 117-136.
12. Cagen SZ, Janoff AS, Bus JS, Gibson JE. Effect of paraquat (methylviologen) on liver function in mice. *J Pharmacol Exp Ther* 1976;198:222-8.
13. Murray RE, Gibson JE. A comparative study of paraquat intoxication in rats, guinea pigs and monkeys. *Exp Mol Pathol* 1972;17: 317-25.
14. Dede EB, Okolonkwo BN, Ngokere AA. Toxicological effect of paraquat on the histology of the stomach, small intestine and testis of male albino rats (*Rattus rattus*). *Port Harcourt Med J* 2007;2:51-5.
15. Wefers H, Sies H. The protection by ascorbate and glutathione against microsomal lipid peroxidation is dependent on vitamin E. *Eur J Biochem* 1988;174:353-7.
16. Wershana KZ. The influence of vitamin C or selenium on paraquat induced toxicity in Guinea Pigs. *Pak J Biol Sci* 2001;4:81-8.
17. Tantcheva LP, Stoeva ES, Galabov AS, Braykova AA, Savov VM, Mileva MM. Effect of vitamin E and vitamin C combination on experimental influenza virus infection. *Exp Clin Pharmacol* 2003;4:259-64.
18. Hassan HM, Fridovich I. Superoxide dismutases: detoxication by a free radical. In: WB Jakoby, ed. *Enzymatic basis of detoxication*. New York: Academic Press; 1980. pp 311-322.
19. Shigeoka S, Ishikawa T, Tamoi M, Miyagawa Y, Takeda T, Yabuta Y, et al. Regulation and function of ascorbate peroxidase isoenzymes. *J Exp Bot* 2002;53: 1305-19.